

# Subclasses of Starlike and Convex Functions Involving Pascal Distribution Series

B. B. Gadekar<sup>1</sup>, R. P. Pathak<sup>2</sup>, S. G. Gajbhiv<sup>3</sup>

<sup>2</sup>Department of Mathematics, School of Science, Sandip University, Nashik, Maharashtra, India,

<sup>2</sup>Department of Mathematics, School of Science, Sandip University, Nashik, Maharashtra, India

<sup>3</sup>Department of Mathematics, MIT Academy of Engineering, Alandi, Pune, Maharashtra, India

Email: [balasahebgadekar@gmail.com](mailto:balasahebgadekar@gmail.com)

The main object of this paper is to establish connections between various subclasses of normalized analytic univalent functions and Pascal distribution series by applying certain convolution operator involving Pascal distribution series.

**Keywords:** Starlike functions, convex functions, uniformly starlike functions, uniformly convex functions, Hadamard product, Pascal distribution series.

## 1. Introduction

Let  $\mathcal{A}$  be the class of functions  $f(z)$  of the form

$$f(z) = z + \sum_{k=2}^{\infty} a_k z^k \quad (1.1)$$

which are analytic in the open unit disk

$$\mathbb{U} = \{z: z \in \mathbb{C} \text{ and } |z| < 1\}$$

As usual, we denote by  $\mathcal{S}$  the subclass of  $\mathcal{A}$  consisting of functions which are normalized by  $f(0) = 0 = f'(0) - 1$  and are also univalent in  $\mathbb{U}$ . Further, let  $\mathcal{T}$  be a subclass of  $\mathcal{A}$  consisting of functions of the form,

$$f(z) = z - \sum_{k=2}^{\infty} a_k z^k, \quad z \in \mathbb{U} \quad (1.2)$$

Also let  $\mathcal{S}^*(\alpha)$  and  $\mathcal{C}(\alpha)$  denote the subclasses of  $\mathcal{A}$  consisting of functions which are, respectively, starlike and convex of order  $\alpha$  ( $0 \leq \alpha < 1$ ) in  $\mathbb{U}$ . Thus, we have (see, for details, [9,13,14,21])

$$\mathcal{S}^*(\alpha) = \{f: f \in \mathcal{A} \text{ and } \operatorname{Re} \left( \frac{zf'(z)}{f(z)} \right) > \alpha, (z \in \mathbb{U}; 0 \leq \alpha < 1)\} \quad (1.3)$$

$$\mathcal{C}(\alpha) = \{f: f \in \mathcal{A} \text{ and } \operatorname{Re} \left( 1 + \frac{zf''(z)}{f'(z)} \right) > \alpha, (z \in \mathbb{U}; 0 \leq \alpha < 1)\} \quad (1.4)$$

where, for convenience,

$$\mathcal{S}^*(0) = \mathcal{S}^*, \quad \mathcal{C}(0) = \mathcal{C}$$

It is a well-established fact that

$$f(z) \in \mathcal{C}(\alpha) \Leftrightarrow zf'(z) \in \mathcal{S}^*(\alpha)$$

Kanas and Wisniowska [15] introduced the class  $\beta$  – UCV which includes geometric aspect in connection with conic domains. The family  $\beta$  – UCV is of special interest for it contains many well-known, as well as new, classes of analytic univalent functions. The class  $\beta$  – UCV map each circular arc contained in the unit disk  $\mathbb{U}$  with a center  $\xi$ ,  $|\xi| \leq \beta$  ( $0 \leq \beta < 1$ ), onto a convex arc. The notion of  $\beta$ -uniformly convex function is simple extension of classical convexity. For various choices of  $\beta$ , these classes reduce to well-known subclasses of uniformly starlike and convex functions. In particular, when  $\beta = 0$ , the center  $\xi$  is the origin and the class  $\beta$ -uniformly convex function reduces to the familiar classes of convex univalent functions  $\mathcal{K}$ . Moreover for  $\beta = 1$  the class reduces to uniformly convex functions UCV introduced by Goodman [12], and studied extensively by Rønning [20]. We recall the following subclass of uniformly convex functions and corresponding subclass of starlike functions due to [20].

For  $-1 < \alpha \leq 1$  and  $\beta \geq 0$ , a function  $f \in \mathcal{A}$  is said to be in the class

i)  $\beta$ -uniformly starlike functions of order  $\alpha$ , denoted by  $S_p(\alpha, \beta)$  if it satisfies analytic criterion

$$\operatorname{Re} \left\{ \frac{zf'(z)}{f(z)} - \alpha \right\} > \beta \left| \frac{zf'(z)}{f(z)} - 1 \right|, \quad z \in \mathbb{U} \quad (1.5)$$

and

ii)  $\beta$ -uniformly convex functions of order  $\alpha$ , denoted by  $UCV(\alpha, \beta)$  if it satisfies analytic criterion

$$\operatorname{Re} \left\{ 1 + \frac{zf''(z)}{f'(z)} - \alpha \right\} > \beta \left| \frac{zf''(z)}{f'(z)} - 1 \right|, \quad z \in \mathbb{U} \quad (1.6)$$

It follows from (1.5) and (1.6) that

$$f \in UCV(\alpha, \beta) \Leftrightarrow zf' \in S_p(\alpha, \beta)$$

We recall the following subclass of uniformly convex functions and corresponding subclass of starlike functions due to [17]. For  $0 \leq \lambda < 1, 0 \leq \alpha < 1$  and  $\beta \geq 0$ , we let  $\mathcal{P}_\lambda(\alpha, \beta)$  be the subclass of  $\mathcal{A}$  consisting of functions of the form (1.1) and satisfying the analytic criterion

$$\operatorname{Re} \left\{ \frac{zf'(z) + \lambda z^2 f''(z)}{(1-\lambda)f(z) + \lambda zf(z)} - \alpha \right\} > \beta \left| \frac{zf'(z) + \lambda z^2 f''(z)}{(1-\lambda)f(z) + \lambda zf(z)} - 1 \right| \quad (1.7)$$

also let  $\mathcal{Q}_\lambda(\alpha, \beta)$  be the subclass of  $\mathcal{S}$  satisfying the analytic criteria

$$\operatorname{Re} \left\{ \frac{\lambda z^2 f'''(z) + (1 + 2\lambda)zf''(z) + f'(z)}{f'(z) + \lambda zf''(z)} - \alpha \right\} > \beta \left| \frac{\lambda z^2 f'''(z) + (1 + \lambda)zf''(z)}{f'(z) + \lambda zf''(z)} \right| \quad (1.8)$$

Also denote  $\mathcal{P}_\lambda^*(\alpha, \beta) = \mathcal{P}_\lambda(\alpha, \beta) \cap \mathcal{T}$  and  $\mathcal{Q}_\lambda^*(\alpha, \beta) = \mathcal{Q}_\lambda(\alpha, \beta) \cap \mathcal{T}$ , the subclasses of  $\mathcal{T}$ . In particular, we note that  $\mathcal{P}_0^*(\alpha, \beta) = \mathcal{TS}_p(\alpha, \beta)$  and  $\mathcal{P}_1^*(\alpha, \beta) = \mathcal{UCT}(\alpha, \beta)$ , the classes of uniformly starlike and uniformly convex were introduced by Bharati et al.[5]. For  $\beta = 0$ , the classes  $\mathcal{TS}_p(\alpha, \beta)$  and  $\mathcal{UCT}(\alpha, \beta)$  respectively, reduces to the classes  $\mathcal{T}^*(\alpha)$  and  $\mathcal{C}(\alpha)$  introduced and studied by Silverman [23]. Further by taking  $\alpha = 0$  we have  $\mathcal{P}_0^*(\beta) = \mathcal{TS}_p(\beta)$  and  $\mathcal{P}_1^*(\beta) = \mathcal{UCT}(\beta)$  introduced and studied by Subramanian et al.[26]. For more interesting developments of some related subclasses of uniformly starlike and uniformly convex functions, the readers may be referred to the works of Frasin et al.[11], Subramanian et al.[25] and Srivastava et al.[24].

A variable  $x$  is said to be Pascal distribution if it takes the values  $0, 1, 2, 3, \dots$  with probabilities  $(1-q)^m, \frac{qm(1-q)^m}{1!}, \frac{q^2m(m+1)(1-q)^m}{2!}, \frac{q^3m(m+1)(m+2)(1-q)^m}{3!}, \dots$  respectively, where  $q$  and  $m$  called parameters, and thus

$$P(X = k) = \binom{k+m-1}{m-1} q^k (1-q)^m, k = 0, 1, 2, 3, \dots \quad (1.9)$$

Recently, El-Deeb et al.[8] introduced a power series whose coefficients are probabilities of Pascal distribution

$$\Psi_q^m(z) = z + \sum_{n=2}^{\infty} \binom{n+m-2}{m-1} \cdot q^{n-1} (1-q)^m z^n, z \in \mathbb{U} \quad (1.10)$$

where  $m \geq 1; 0 \leq q \leq 1$  and we note that, by ratio test the radius of convergence of above series is infinity. We also define the series

$$\Phi_q^m(z) = 2z - \Psi_q^m(z) = z - \sum_{n=2}^{\infty} \binom{n+m-2}{m-1} \cdot q^{n-1} (1-q)^m z^n, z \in \mathbb{U} \quad (1.11)$$

Now, we considered the linear operator  $\mathcal{J}_q^m(z): \mathcal{A} \rightarrow \mathcal{A}$  defined by Hadamard product (convolution) as

$$\mathcal{J}_q^m(z) = \Psi_q^m(z) * f(z) = z + \sum_{n=2}^{\infty} \binom{n+m-2}{m-1} \cdot q^{n-1} (1-q)^m z^n, z \in \mathbb{U} \quad (1.12)$$

To establish our main results, we need the following Lemmas.  
 Lemma 1.1. [17] A function  $f$  of the form (1.2) is in  $\mathcal{P}_\lambda^*(\alpha, \beta)$  if and only if it satisfies

$$\sum_{n=2}^{\infty} (1 + n\lambda - \lambda)[n(1 + \beta) - (\alpha + \beta)]|a_n| \leq 1 - \alpha \quad (1.13)$$

Lemma 1.2. [17] A function  $f$  of the form (1.2) is in  $\mathcal{Q}_\lambda^*(\alpha, \beta)$  if and only if it satisfies

$$\sum_{n=2}^{\infty} n(1 + n\lambda - \lambda)[n(1 + \beta) - (\alpha + \beta)]|a_n| \leq 1 - \alpha \quad (1.14)$$

Lemma 1.3. [7] If  $f \in \mathcal{R}^*(C, D)$  is of the form (1.1) then

$$|a_n| = \frac{(C - D)|v|}{n}, n \in \mathbb{N} \setminus \{1\} \quad (1.15)$$

Motivated by several earlier results on connections between various subclasses of analytic and univalent functions by using hypergeometric functions [6, 17, 18, 19, 22] and by the recent investigation in generalized Bessel functions and Struve functions [1, 2, 3, 10, 24, 27, 28] in the present paper we establish a number of connections between the classes  $\mathcal{P}_\lambda^*(\alpha, \beta)$  and  $\mathcal{Q}_\lambda^*(\alpha, \beta)$  by applying the convolution operator given by (1.12). We also determine conditions for the integral operator  $\mathcal{H}(m, z) = \int_0^z \frac{t^{mf(t)}}{t} dt$  belonging to the class  $\mathcal{Q}_\lambda^*(\alpha, \beta)$ .

## 2. Coefficient Estimates

Theorem 2.1. If  $m \geq 1$ , then  $\Phi_q^m(z)$  is in  $\mathcal{P}_\lambda^*(\alpha, \beta)$  if and only if

$$(1 + \beta)\lambda \frac{q^2 m(m + 1)}{(1 - q)^2} + [(1 + \beta)(1 + 2\lambda) - \lambda(\alpha + \beta)] \frac{qm}{(1 - q)}$$

Proof. Since

$$\Phi_q^m(z) = z - \sum_{n=2}^{\infty} \binom{n + m - 2}{m - 1} \cdot q^{n-1} (1 - q)^m a_n z^n$$

in view of Lemma 1.1, it is sufficient to show that

$$\sum_{n=2}^{\infty} (1 + n\lambda - \lambda)[n(1 + \beta) - (\alpha + \beta)] \binom{n + m - 2}{m - 1} \cdot q^{n-1} (1 - q)^m \leq 1 - \alpha \quad (2.2)$$

In the proofs of all our results we will use the relation

$$\sum_{n=0}^{\infty} \binom{n + m - 2}{m - 1} q^n = \frac{1}{(1 - q)^m}, \quad (0 \leq q \leq 1)$$

and the corresponding ones obtained by replacing the value of  $m$  with  $m - 1$ ,  $m + 1$ , and  $m +$

2.

Writing  $n^2 = (n-2)(n-1) + 3(n-1) + 1$  and  $n = (n-1) + 1$  in (2.2)

$$\begin{aligned}
 & \sum_{n=2}^{\infty} (1+n\lambda-\lambda)[n(1+\beta)-(\alpha+\beta)] \binom{n+m-2}{m-1} \cdot q^{n-1}(1-q)^m \\
 &= (1+\beta)\lambda \sum_{n=2}^{\infty} [(n-2)(n-1) + 3(n-1) + 1] \binom{n+m-2}{m-1} \cdot q^{n-1}(1-q)^m \\
 &+ [(1+\beta)(1-\lambda) - \lambda(\alpha+\beta)] \sum_{n=2}^{\infty} [(n-1) + 1] \binom{n+m-2}{m-1} \cdot q^{n-1}(1-q)^m \\
 &+ (1-\alpha) \sum_{n=2}^{\infty} \binom{n+m-2}{m-1} \cdot q^{n-1}(1-q)^m \\
 &= (1+\beta)\lambda q^2 m(m+1)(1-q)^m \sum_{n=0}^{\infty} \binom{n+m+1}{m+1} \cdot q^n \\
 &+ [(1+\beta)(1-\lambda) - \lambda(\alpha+\beta)] qm(1-q)^m \sum_{n=0}^{\infty} \binom{n+m}{m} \cdot q^n \\
 &+ (1-\alpha)(1-q)^m \sum_{n=0}^{\infty} \binom{n+m-1}{m-1} \cdot q^{n-1} \\
 &= (1+\beta)\lambda q^2 m(m+1)(1-q)^m \frac{1}{(1-q)^{m+2}} + [(1+\beta)(1-\lambda) - \lambda(\alpha+\beta)] qm \\
 &\cdot (1-q)^m \frac{1}{(1-q)^{m+1}} + (1-\alpha)(1-q)^m \left[ \frac{1}{(1-q)^m} - 1 \right] \\
 &= (1+\beta)\lambda q^2 m(m+1) \frac{1}{(1-q)^2} + [(1+\beta)(1+2\lambda) - \lambda(\alpha+\beta)] \frac{qm}{(1-q)} \\
 &\cdot (1-\alpha)[1 - (1-q)^m]
 \end{aligned}$$

But this last expression is bounded above by  $1-\alpha$  if and only if (2.1) holds. Theorem 2.2. If  $m \geq 1$ , then  $\phi_q^m(z)$  is in  $\mathcal{Q}_\lambda^*(\alpha, \beta)$  if and only if

$$= (1+\beta)\lambda \frac{q^3 m(m+1)(m+2)}{(1-q)^3} + [(1+\beta)(1+5\lambda) - \lambda(\alpha+\beta)] \frac{q^2 m(m+1)}{(1-q)^2}$$

Proof. The proof of Theorem 2.2 is lines similar to the proof of Theorem 2.1, so we omitted the proof.

By taking  $\lambda = 0$  and  $\lambda = 1$  we state the following Corollaries. Corollary 2.1. If  $m \geq 1$  then necessary and sufficient condition for  $\phi_q^m(z) \in \mathcal{P}_0^*(\alpha, \beta) \equiv \mathcal{TS}_P(\alpha, \beta)$  is

$$(1 + \beta) \frac{qm}{(1 - q)} + (1 - \alpha)[1 - (1 - q)^m] \leq 1 - \alpha \quad (2.4)$$

Corollary 2.2. If  $m \geq 1$  then necessary and sufficient condition for  $\phi_q^m(z) \in \mathcal{P}_1^*(\alpha, \beta) \equiv \mathcal{UCT}(\alpha, \beta)$  is

$$(1 + \beta) \frac{q^2 m(m + 1)}{(1 - q)^2} + [3(1 + \beta) - (\alpha + \beta)] \frac{qm}{(1 - q)}$$

### 3. Inclusion Properties

Theorem 3.1. If  $f \in \mathcal{R}^r(C, D)$  is of the form (1.1) and the operator  $\mathcal{J}_q^m(z)$ , defined by (1.12) in the class  $\mathcal{P}_\lambda^*(\alpha, \beta)$ , if

$$(C - D)|v|(1 - q)^m [(1 + \beta)\lambda \left( \frac{qm}{(1 - q)} \right) + [(1 + \beta) - \lambda(\alpha + \beta)](1 - (1 - q)^m)]$$

Proof. In view of Lemma 1.1, it is sufficient to show that

$$\sum_{n=2}^{\infty} (1 + n\lambda - \lambda)[n(1 + \beta) - (\alpha + \beta)] \binom{n + m - 2}{m - 1} \cdot q^{n-1} |a_n| (1 - q)^m \leq 1 - \alpha \quad (3.2)$$

Since  $f \in \mathcal{R}^r(C, D)$ , then by Lemma 1.3 we have

$$|a_n| = \frac{(C - D)|v|}{n}.$$

Thus, we have

$$\begin{aligned} &= \sum_{n=2}^{\infty} (1 + n\lambda - \lambda)[n(1 + \beta) - (\alpha + \beta)] \binom{n + m - 2}{m - 1} \cdot q^{n-1} |a_n| (1 - q)^m \\ &\leq \sum_{n=2}^{\infty} \frac{(C - D)|v|}{n} (1 + n\lambda - \lambda)[n(1 + \beta) - (\alpha + \beta)] \binom{n + m - 2}{m - 1} \cdot q^{n-1} (1 - q)^m \\ &= (C - D)|v| \left[ \sum_{n=2}^{\infty} (1 + n\lambda - \lambda)[n(1 + \beta) - (\alpha + \beta)] \binom{n + m - 2}{m - 1} \cdot q^{n-1} (1 - q)^m \right] \\ &= (C - D)|v|(1 - q)^m \left[ ((1 + \beta)\lambda \sum_{n=2}^{\infty} \binom{n + m - 2}{m - 1} \cdot q^{n-1} + [(1 + \beta) - (\alpha + \beta)] \right. \\ &\quad \times \sum_{n=2}^{\infty} \binom{n + m - 2}{m - 1} q^{n-1} - (1 - \lambda)(\alpha + \beta) \sum_{n=2}^{\infty} \frac{1}{n} \binom{n + m - 2}{m - 1} \cdot q^{n-1} \left. \right] \end{aligned}$$

writing  $n = (n - 1) + 1$ , upon simple computation

$$\begin{aligned}
 &= (C - D)|v|(1 - q)^m [(1 + \beta)\lambda (qm \sum_{n=0}^{\infty} \binom{n+m}{m} \cdot q^n + \sum_{n=0}^{\infty} \binom{n+m-1}{m-1} \cdot q^n - 1) \\
 &+ [(1 + \beta)(1 - \lambda) - \lambda(\alpha + \beta)] (\sum_{n=0}^{\infty} \binom{n+m-1}{m-1} \cdot q^n - 1) \\
 &- \frac{(1 - \lambda)(\alpha + \beta)}{q(m-1)} (\sum_{n=0}^{\infty} \binom{n+m-2}{m-2} \cdot q^n - 1 - (m-1)q)] \\
 &= (C - D)|v|(1 - q)^m [(1 + \beta)\lambda (\frac{qm}{(1-q)} + 1 - (1 - q)^m) + [(1 + \beta)(1 - \lambda) - \lambda(\alpha + \beta)] \\
 &\times (1 - (1 - q)^m) - \frac{(1 - \lambda)(\alpha + \beta)}{q(m-1)} ((1 - q) - (1 - q)^m - q(m-1)(1 - q)^m)] \\
 &= (C - D)|v|(1 - q)^m [(1 + \beta)\lambda (\frac{qm}{(1-q)}) + [(1 + \beta) - \lambda(\alpha + \beta)] \\
 &\times (1 - (1 - q)^m) - \frac{(1 - \lambda)(\alpha + \beta)}{q(m-1)} ((1 - q) - (1 - q)^m - q(m-1)(1 - q)^m)]
 \end{aligned}$$

Theorem 3.2. If  $f \in \mathcal{R}^{\tau}(C, D)$  is of the form (1.1) and the operator  $\mathcal{J}_q^m(z)$ , defined by (1.12) in the class  $\mathcal{Q}_{\lambda}^*(\alpha, \beta)$ , if

$$(C - D)|v| [(1 + \beta)\lambda \frac{q^2 m(m+1)}{(1-q)^2} + [(1 + \beta)(1 + 2\lambda) - \lambda(\alpha + \beta)] \frac{qm}{(1-q)}]$$

Proof. In view of Lemma 1.2, it is sufficient to show that

$$\sum_{n=2}^{\infty} n(1 + n\lambda - \lambda)[n(1 + \beta) - (\alpha + \beta)] \binom{n+m-2}{m-1} \cdot q^{n-1} |a_n| (1 - q)^m \leq 1 - \alpha \quad (3.4)$$

Since  $f \in \mathcal{R}^{\tau}(C, D)$ , then by Lemma 1.3 we have

$$|a_n| = \frac{(C - D)|v|}{n}$$

Thus we have

$$\begin{aligned}
 &\Rightarrow \sum_{n=2}^{\infty} \frac{(C - D)|v|}{n} n(1 + n\lambda - \lambda)[n(1 + \beta) - (\alpha + \beta)] \binom{n+m-2}{m-1} \cdot q^{n-1} (1 - q)^m \\
 &\Rightarrow \leq (C - D)|v|(1 - q)^m \sum_{n=2}^{\infty} (1 + n\lambda - \lambda)[n(1 + \beta) - (\alpha + \beta)] \binom{n+m-2}{m-1} \cdot q^{n-1}
 \end{aligned}$$

Further, proceeding as in Theorem 2.1, we get

$$\begin{aligned}
 &(C - D)|v|(1 + \beta)\lambda \frac{q^2 m(m+1)}{(1-q)^2} + [(1 + \beta)(1 + 2\lambda) - \lambda(\alpha + \beta)] \frac{qm}{(1-q)} \\
 &+ (1 - \alpha)[1 - (1 - q)^m] \leq 1 - \alpha
 \end{aligned}$$

Corollary 3.1. Let  $m \geq 1$ . If  $f \in \mathcal{R}^{\tau}(C, D)$  is of the form (1.1) and the operator  $\mathcal{J}_q^m(z)$ , defined by (1.12) in the class  $\mathcal{P}_0^*(\alpha, \beta) \equiv \mathcal{TS}_p(\alpha, \beta)$ , if

$$\Rightarrow (C - D)|v|(1 - q)^m [(1 + \beta)(1 - (1 - q)^m) \frac{(\alpha + \beta)}{q(m - 1)}]$$

Corollary 3.2. Let  $m \geq 1$ . If  $f \in \mathcal{R}^{\tau}(C, D)$  is of the form (1.1) and the operator  $\mathcal{J}_q^m(z)$ , defined by (1.12) in the class  $\mathcal{P}_0^*(\alpha, \beta) \equiv \mathcal{TS}_p(\alpha, \beta)$ , if

$$(C - D)|v| [(1 + \beta) \frac{q^2 m(m + 1)}{(1 - q)^2} + [3(1 + \beta) - (\alpha + \beta)] \frac{qm}{(1 - q)}]$$

#### 4. An Integral Operator

In this section we introduce an integral operator  $\mathcal{H}(m, z)$  as follows:

$$\mathcal{H}(m, z) = \int_0^z \frac{\mathcal{J}_q^m f(t)}{t} dt \quad (4.1)$$

and we obtain a necessary and sufficient condition for  $\mathcal{H}(m, z)$  belonging to the class  $\mathcal{Q}_{\lambda}^*(\alpha, \beta)$ .

Theorem 4.1. If  $m \geq 1$ , then the integral  $\mathcal{H}(m, z)$  defined by (4.1) is in the class  $\mathcal{Q}_{\lambda}^*(\alpha, \beta)$ , if and only if inequality (2.1) is satisfied.

Proof. Since

$$\mathcal{H}(z) = z + \sum_{n=2}^{\infty} \binom{n+m-2}{m-1} \cdot q^{n-1} (1-q)^m \frac{z^n}{n}$$

in view of Lemma (1.2), we need only to show that

$$\sum_{n=2}^{\infty} n(1+n\lambda-\lambda)[n(1+\beta)-(\alpha+\beta)] \times \frac{1}{n} \binom{n+m-2}{m-1} \cdot q^{n-1} (1-q)^m \leq 1-\alpha$$

or, equivalently

$$\sum_{n=2}^{\infty} (1+n\lambda-\lambda)[n(1+\beta)-(\alpha+\beta)] \binom{n+m-2}{m-1} \cdot q^{n-1} (1-q)^m \leq 1-\alpha$$

The remaining part of the proof of Theorem 4.1 is similar to that of Theorem 2.1, and so we omit the details.

#### 5. Conclusion

In this investigation, we obtain sufficient condition and inclusion results for function  $f \in \mathcal{A}$  to be in the classes of starlike and convex functions  $\mathcal{P}_{\lambda}^*(\alpha, \beta)$  and  $\mathcal{Q}_{\lambda}^*(\alpha, \beta)$  associated with Pascal distribution series in the open unit disc  $\mathbb{U}$ . Further, we derive the result of integral operator



related to Pascal distribution series.

## References

1. Baricz, A., Generalized Bessel functions of the first kind, Lecture Notes in Math., 1994, SpringerVerlag(2010).
2. Baricz, A., Geometric properties of generalized Bessel functions of complex order, *Mathematica(Cluj)*,48,13-18,(2006).
3. Baricz, A., Geometric properties of generalized Bessel functions, *Publ. Math. Debrecen*, 73, 155-178,(2008).
4. de Branges, L., A proof of the Bieberbach conjecture, *Acta Math.*, 154, 137-152, (1985).
5. Bharati, R., Parvatham, R. and Swaminathan, A., On subclasses of uniformly convex functions and corresponding class of starlike functions, *Tamkang J. Math.*, 26, no.1, 17-32, (1997).
6. Cho, N. E., Woo S. Y. and Owa, S., Uniform convexity properties for hypergeometric functions, *Fract.Cal.Appl.Anal.*,5,no.3,303-313,(2002).
7. Dixit, K. K. and Pal, S. K., On a class of univalent functions related to complex order, *Indian J. Pure.Appl.Math.*,26,no.9,889-896,(1995).
8. El-Deeb, S. M., Bulboacă T. and Dziok, J., Pascal distribution series connected with certain subclasses of univalent functions, *Kyungpook Math. J.*, 59, 301-314, (2019).
9. Duren, P. L., *Univalent Functions*, Grundlehren der Mathematischen Wissenschaften, Band 259, Springer-Verlag, New York, Berlin, Heidelberg and Tokyo, (1983).
10. Deniz, E., Orhan H. and Srivastava, H. M., Some sufficient conditions for univalence of certain families of integral operators involving generalized Bessel functions, *Taiwanese J. Math.*, 15, 883917,(2011).
11. Frasin, B. A., Al-Hawary, T. and Yousef, F., Necessary and sufficient conditions for hypergeometric functions to be in a subclass of analytic functions, *Afr. Mat.*, 30, no.1-2, 223-230, (2019).
12. Goodman, A. W., On uniformly convex functions, *Ann. Polon. Math.*, 56, 87-92, 1991.
13. Kamble P. N., Shrigan M. G. and Srivastava H. M., A novel subclass of univalent functions involving operators of fractional calculus, *Int. J. Appl. Math.*, 30, no. 6, 501-514, (2017).
14. Kamble P. N., Shrigan M. G., Coefficient estimates for a subclass of Bi-univalent functions defined by Sălăgean Type  $q$ -Calculus operator, 48, 677-688, (2018).
15. Kanas, S. and Wisniowska, Conic regions and  $k$ -uniform convexity, *Comput. Appl. Math.*, 105,327-336,1999.
16. Kanas, S. and Srivastava, H. M., Linear operators associated with  $k$ -uniformly convex functions, *Integral Transform. Spec. Funct.*, 9, no.2, 121-132, (2000).
17. Murugusundaramoorthy, G., Subclasses of starlike and convex functions involving Poisson distribution series, *Afr. Mat.*, 28:1357-1366, (2017).
18. Merkes, E. P. and Scott W. P., Starlike hypergeometric functions, *Proc. Amer. Math. Soc.*, 12, 885-888,(1961).
19. Owa, S. and Srivastava, H. M., Univalent and starlike generalized hypergeometric functions, *Canad. J. Math.*, 39, 1057-1077, (1987).
20. Rønning, F., Uniformly convex functions and a corresponding class of starlike functions, *Proc. Amer.Math.Soc.*,118,189-196,1993.
21. Shrigan M. G., Second Hankel determinant for bi-univalent functions associated with  $q$ -differential operator, *J. Sib. Fed. Univ. Math. Phys.*, 15 (5) (2022), 663-671.
22. Shrigan, M. G., Yalçın S. and Altınkaya, S., Unified approach to starlike and convex functions involving Poisson distribution series, *Bul. Acad. Ştiinţe Repub. Moldova Mat.*, 97, no.3, 11-20,

- (2021).
23. Silverman, H., Univalent functions with negative coefficients, Proc. Amer. Math. Soc., 51, 109-116,(1975).
  24. Srivastava, H. M., Murugusundaramoorthy, G. and Janani, T., Uniformly Starlike Functions and Uniformly Convex Functions Associated with the Struve Function, J. Appl. Computat. Math., 3:6,(2014).
  25. Subramanian, K. G., Sudharsan, T. V., Balasubrahmanyam, P. and Silverman, H., Classes of uniformly starlike functions, Publ. Math. Debrecen., 53, no. 3-4, 309-315, (1998).
  26. Subramanian, K. G., Murugusundaramoorthy, G., Balasubrahmanyam, P. and Silverman, H., Subclasses of uniformly convex and uniformly starlike functions, Math. Japonica, 42, no.3, 517-522,(1995).
  27. Yagmur, N. and Orhan, H., Starlikeness and Convexity of Generalized Struve Functions, Abstr. Appl. Anal., 2013, Article ID954513, 6pp.
  28. Zhang, S. and Jin, J. Computation of Special Functions, A Wiley-Interscience Publication, John Wiley and Sons, Inc., New York, (1996).