

Contra M -Open Maps and Homeomorphisms in Pythagorean Fuzzy Topological Spaces

B. Vijayalakshmi, M. Ramalakshmi, A. Vadivel, G. Saravanakumar

1 Introduction

Considering the imprecision in decision-making, Zadeh [39] introduced the idea of fuzzy set which has a membership function, μ that assigns to each element of the universe of discourse, a number from the unit interval $[0,1]$ to indicate the degree of belongingness to the set under consideration. The notion of fuzzy sets generalizes classical sets theory by allowing intermediate situations between the whole and nothing. In a fuzzy set, a membership function is defined to describe the degree of membership of an element to a class. The membership value ranges from 0 to 1, where 0 shows that the element does not belong to a class, 1 means belongs, and other values indicate the degree of membership to a class. For fuzzy sets, the membership function replaced the characteristic function in crisp sets. The concept of fuzzy set theory seems to be inconclusive because of the exclusion of nonmembership function and the disregard for the possibility of hesitation margin. Atanassov critically studied these shortcomings and proposed a concept called intuitionistic fuzzy sets (IFS s) [1, 2, 4, 5]. The construct (that is, IFS s) incorporates both membership function, μ and nonmembership function, ν with hesitation margin, π (that is, neither membership nor non-membership functions), such that $\mu + \nu \leq 1$ and $\mu + \nu + \pi = 1$. Atanassov [3] introduced intuitionistic fuzzy sets of second type ($IFSST$) with the property that the sum of the square of the membership and non-membership degrees is less than or equal to one. This concept generalizes IFS s in a way. The notion of IFS s provides a flexible framework to elaborate uncertainty and vagueness. The idea of IFS seems to be resourceful in modelling many real-life situations like medical diagnosis [7, 8, 13, 30, 31], career determination [11], selection process [12], and multi-criteria decision-making [16, 17, 18], among others.

There are situations where $\mu + \nu \geq 1$ unlike the cases capture in IFS s. This limitation in IFS naturally led to a construct, called Pythagorean fuzzy sets ($pfss$). Pythagorean fuzzy set (pfs) proposed in [36, 37, 38] is a new tool to deal with vagueness considering the membership grade, μ and non-membership grade, ν satisfying the conditions $\mu + \nu \leq 1$ or $\mu + \nu \geq 1$, and also, it follows that $\mu^2 + \nu^2 + \pi^2 = 1$, where π is the Pythagorean fuzzy set index. In fact, the origin of Pythagorean fuzzy sets emanated from $IFSST$ earlier studied in the literature. As a generalized set, PFS has close relationship with IFS . The construct of PFS s can be used to characterize uncertain information more sufficiently and accurately than IFS . Garg [15] presented an improved score function for the ranking order of interval-valued Pythagorean fuzzy sets ($IVPFS$ s). Based on it, a Pythagorean fuzzy technique for order of preference by similarity to ideal solution ($TOPSIS$) method by taking the preferences of the experts in the form of interval-valued Pythagorean fuzzy decision matrices was discussed. Other explorations of the theory of PFS s can be found in [6, 9, 14, 19, 20, 26, 27]. Saha [29] defined δ -open sets in topological spaces. Vadivel et al. [34] introduced δ -open sets in a neutrosophic topological space. The notion of M -open sets in topological spaces were introduced by El-Maghrabi and Al-Juhani [24] in 2011 and studied some of their properties. The class of sets namely, M -open sets are playing more important role in topological spaces, because of their applications in various fields of Mathematics and other real fields. Recently, Jeeva et al. [21, 22, 23] introduced neutrosophic soft M -open sets in neutrosophic topological spaces and developed the concepts of neutrosophic soft M -Continuity and M -Irresolute maps.

The concept of contra continuous function in general topology was introduced by Dontchev [10] in 1996. Vadivel et al. [33] introduced generalized fuzzy contra e -continuous functions in fuzzy topological spaces. Recently, Revathi et al. [?] developed the neutrosophic soft contra e -continuity and e -irresolute maps in neutrosophic soft topological spaces.

The focus of this article is to introduce the idea of Pythagorean fuzzy contra M -open and Pythagorean fuzzy contra M -closed mappings in Pythagorean fuzzy topological spaces. Also the work is extended to Pythagorean fuzzy contra M -homeomorphism and Pythagorean fuzzy contra M C -homeomorphism in Pythagorean fuzzy topological spaces and some of their basic properties are discussed.

2 Preliminaries

We recall some basic notions of fuzzy sets, IFS s and pf s.

Definition 2.1 [39] Let X be a nonempty set. A fuzzy set A in X is characterized by a membership function $\mu_A: X \rightarrow [0,1]$. That is:

$$\mu_A(x) = \begin{cases} 1, & \text{if } x \in A \\ 0, & \text{if } x \notin A \\ (0,1) & \text{if } x \text{ is partly in } A. \end{cases}$$

Alternatively, a fuzzy set A in X is an object having the form $A = \{ \langle x, \mu_A(x) \rangle \mid x \in X \}$ or $A = \{ \langle \frac{\mu_A(x)}{x} \rangle \mid x \in X \}$, where the function $\mu_A(x): X \rightarrow [0,1]$ defines the degree of membership of the element, $x \in X$.

The closer the membership value $\mu_A(x)$ to 1, the more x belongs to A , where the grades 1 and 0 represent full membership and full nonmembership. Fuzzy set is a collection of objects with graded membership, that is, having degree of membership. Fuzzy set is an extension of the classical notion of set. In classical set theory, the membership of elements in a set is assessed in a binary terms according to a bivalent condition; an element either belongs or does not belong to the set. Classical bivalent sets are in fuzzy set theory called crisp sets. Fuzzy sets are generalized classical sets, since the indicator function of classical sets is special cases of the membership functions of fuzzy sets, if the latter only take values 0 or 1. Fuzzy sets theory permits the gradual assessment of the membership of element in a set; this is described with the aid of a membership function valued in the real unit interval $[0,1]$.

Let us consider two examples:

(i) all employees of XYZ who are over 1.8m in height; (ii) all employees of XYZ who are tall. The first example is a classical set with a universe (all XYZ employees) and a membership rule that divides the universe into members (those over 1.8m) and nonmembers. The second example is a fuzzy set, because some employees are definitely in the set and some are definitely not in the set, but some are borderline.

This distinction between the ins, the outs, and the borderline is made more exact by the membership function, μ . If we return to our second example and let A represent the fuzzy set of all tall employees and x represent a member of the universe X (i.e. all employees), then $\mu_A(x)$ would be $\mu_A(x) = 1$ if x is definitely tall or $\mu_A(x) = 0$ if x is definitely not tall or $0 < \mu_A(x) < 1$ for borderline cases.

Definition 2.2 [1, 2, 4, 5] Let a nonempty set X be fixed. An IFS A in X is an object having the form: $A = \{ \langle x, \mu_A(x), \nu_A(x) \rangle \mid x \in X \}$ or $A = \{ \langle \frac{\mu_A(x), \nu_A(x)}{x} \rangle \mid x \in X \}$, where the

functions $\mu_A(x): X \rightarrow [0,1]$ and $\nu_A(x): X \rightarrow [0,1]$ define the degree of membership and the degree of nonmembership, respectively, of the element $x \in X$ to A , which is a subset of X , and for every $x \in X$: $0 \leq \mu_A(x) + \nu_A(x) \leq 1$. For each A in X : $\pi_A(x) = 1 - \mu_A(x) - \nu_A(x)$ is the intuitionistic fuzzy set index or hesitation margin of x in X . The hesitation margin $\pi_A(x)$ is the degree of nondeterminacy of $x \in X$ to the set A and $\pi_A(x) \in [0,1]$. The hesitation margin is the function that expresses lack of knowledge of whether $x \in X$ or $x \notin X$. Thus: $\mu_A(x) + \nu_A(x) + \pi_A(x) = 1$.

Example 2.1 Let $X = \{x, y, z\}$ be a fixed universe of discourse and $A = \{ \langle \frac{0.6, 0.1}{x} \rangle, \langle \frac{0.5, 0.3}{y} \rangle, \langle \frac{0.2, 0.3}{z} \rangle \}$ be the intuitionistic fuzzy set in X . The hesitation margins of the elements x, y, z to A are as follows: $\pi_A(x) = 0.3, \pi_A(y) = 0.1$ and $\pi_A(z) = 0.2$.

Definition 2.3 [36, 37, 38] Let X be a universal set. Then, a Pythagorean fuzzy set A , which is a set of ordered pairs over X , is defined by the following: $A = \{ \langle x, \mu_A(x), \nu_A(x) \rangle | x \in X \}$ or $A = \{ \langle \frac{\mu_A(x), \nu_A(x)}{x} \rangle | x \in X \}$, where the functions $\mu_A(x): X \rightarrow [0,1]$ and $\nu_A(x): X \rightarrow [0,1]$ define the degree of membership and the degree of nonmembership, respectively, of the element $x \in X$ to A , which is a subset of X , and for every $x \in X$, $0 \leq (\mu_A(x))^2 + (\nu_A(x))^2 \leq 1$. Supposing $(\mu_A(x))^2 + (\nu_A(x))^2 \leq 1$, then there is a degree of indeterminacy of $x \in X$ to A defined by $\pi_A(x) = \sqrt{1 - [(\mu_A(x))^2 + (\nu_A(x))^2]}$ and $\pi_A(x) \in [0,1]$. In what follows, $(\mu_A(x))^2 + (\nu_A(x))^2 + (\pi_A(x))^2 = 1$. Otherwise, $\pi_A(x) = 0$ whenever $(\mu_A(x))^2 + (\nu_A(x))^2 = 1$. We denote the set of all PFS's over X by $pfs(X)$.

Definition 2.4 [38] Let A and B be pfs's of the forms $A = \{ \langle a, \lambda_A(a), \mu_A(a) \rangle | a \in X \}$ and $B = \{ \langle a, \lambda_B(a), \mu_B(a) \rangle | a \in X \}$. Then [(i)]

1. $A \subseteq B$ if and only if $\lambda_A(a) \leq \lambda_B(a)$ and $\mu_A(a) \geq \mu_B(a)$ for all $a \in X$.
2. $A = B$ if and only if $A \subseteq B$ and $B \subseteq A$.
3. $\bar{A} = \{ \langle a, \mu_A(a), \lambda_A(a) \rangle | a \in X \}$.
4. $A \cap B = \{ \langle a, \lambda_A(a) \wedge \lambda_B(a), \mu_A(a) \vee \mu_B(a) \rangle | a \in X \}$.
5. $A \cup B = \{ \langle a, \lambda_A(a) \vee \lambda_B(a), \mu_A(a) \wedge \mu_B(a) \rangle | a \in X \}$.
6. $0_X = \{ \langle a, 0, 1 \rangle | a \in X \}$ and $1_X = \{ \langle a, 1, 0 \rangle | a \in X \}$.
7. $\bar{1} = 0$ and $\bar{0} = 1$.

Definition 2.5 [25] An Pythagorean fuzzy topology by subsets of a non-empty set X is a family τ of pfs's satisfying the following axioms. [(i)]

1. $\phi, X \in \tau$.
2. $G_1 \cap G_2 \in \tau$ for every $G_1, G_2 \in \tau$ and
3. $\bigcup G_i \in \tau$ for any arbitrary family $\{G_i | i \in j\} \subseteq \tau$. The pair (X, τ) is called an Pythagorean fuzzy topological space (*pfts* in short) and any pfs G in τ is called an Pythagorean fuzzy open set (*pfos* in short) in X . The complement \bar{A} of an Pythagorean fuzzy open set A in an *pfts* (X, τ) is called an Pythagorean fuzzy closed set (*pfc*s in short).

Definition 2.6 [25] Let (X, τ) be an *pfts* and $A = \{ \langle a, \lambda_A(a), \mu_A(a) \rangle \mid a \in X \}$ be an *pfs* in X . Then the interior and the closure of A are denoted by $pfint(A)$ and $pfcl(A)$ and are defined as follows: $pfcl(A) = \cap \{K \mid K \text{ is an } pfcs \text{ and } A \subseteq K\}$ and $pfint(A) = \cup \{G \mid G \text{ is an } pfos \text{ and } G \subseteq A\}$. Also, it can be established that $pfcl(A)$ is an *pfcs* and $pfint(A)$ is an *pfos*, A is an *pfcs* if and only if $pfcl(A) = A$ and A is an *pfos* if and only if $pfint(A) = A$. We say that A is *pf-dense* if $pfcl(A) = X$.

Lemma 2.1 [32] For any Pythagorean fuzzy set A in (X, τ) , we have $X - pfint(A) = pfcl(X - A)$ and $X - pfcl(A) = pfint(X - A)$.

Definition 2.7 [32] Let (X, τ) be an *pfts* and A be an *pfs*. Then A is said to be an Pythagorean fuzzy (i) regular open set (*pfros* in short) if $A = pfint(pfcl(A))$. (ii) regular closed set (*pfrcs* in short) if $A = pfcl(pfint(A))$. By Lemma 2.1, it follows that A is an *pfros* iff A is an *pfrcs*.

Definition 2.8 [35] Let (X_1, Γ_P) (or X_1) be an *pfts* and $A = \{ \langle a, \lambda_A(a), \mu_A(a) \rangle \mid a \in X_1 \}$ be an *pfs* in X_1 . Then the (i) $pf\delta$ -interior of A are denoted by $pf\delta int(A)$ and are defined as follows. $pf\delta int(A) = \cup \{G \mid G \text{ is an } pfros \text{ and } G \subseteq A\}$. (ii) $pf\delta$ -closure of A are denoted by $pf\delta cl(A)$ and are defined as follows. $pf\delta cl(A) = \cap \{K \mid K \text{ is an } pfrcs \text{ and } A \subseteq K\}$.

Definition 2.9 [35] Let (X_1, Γ_P) be an *pfts* and $A = \{ \langle a, \lambda_A(a), \mu_A(a) \rangle \mid a \in X_1 \}$ be an *pfs* in X_1 . A set A is said to be *pf* [(i)]

1. δ -open set (briefly, *pf* δ os) if $A = pf\delta int(A)$,
2. δ -pre open set (briefly, *pf* δ Pos) if $A \subseteq pfint(pf\delta cl(A))$,
3. δ -semi open set (briefly, *pf* δ Sos) if $A \subseteq pfcl(pf\delta int(A))$,
4. *e* open set (briefly, *pfeos*) if $A \subseteq pfcl(pf\delta int(A)) \cup pfint(pf\delta cl(A))$,
5. δ (resp. δ -pre, δ -semi and *e*) dense if $pf\delta cl(A)$ (resp. *pf* δ Pcl(A), *pf* δ Scl(A) and *pfecl*(A)) = X_1 .

The complement of an *pf* δ os (resp. *pf* δ Pos, *pf* δ Sos and *pfeos*) is called an *pf* δ (resp. *pf* δ P, *pf* δ S and *pfe*) closed set (briefly, *pf* δ cs (resp. *pf* δ Pcs, *pf* δ Scs and *pfe*cs)) in X_1 .

The family of all *pf* δ os (resp. *pf* δ cs, *pf* δ Pos, *pf* δ Pcs, *pf* δ Sos, *pf* δ Scs, *pfeos* and *pfe*cs) of X_1 is denoted by *pf* δ OS(X_1), (resp. *pf* δ CS(X_1), *pf* δ POS(X_1), *pf* δ Pcs(X_1), *pf* δ SOS(X_1), *pf* δ SCS(X_1), *pfeOS*(X_1) and *pfeCS*(X_1)).

Definition 2.10 [35] Let (X, τ) be an *pfts* and $A = \{ \langle a, \lambda_A(a), \mu_A(a) \rangle \mid a \in X_1 \}$ be an *pfs* in X_1 . Then the (i) *pf* δ -pre (resp. *pf* δ -semi and *pfe*)-interior of A are denoted by *pf* δ Pint(A) (resp. *pf* δ Sint(A) and *pfeint*(A)) and are defined as follows: *pf* δ Pint(A) (resp. *pf* δ Sint(A) and *pfeint*(A)) = $\cup \{G \mid G \text{ is a } pf\delta Pos \text{ (resp. } pf\delta Sos \text{ and } pfeos) \text{ and } G \subseteq A\}$.

$G \subseteq A$ }, (ii) $pf\delta$ -pre (resp. $pf\delta$ -semi and pfe)-closure of A are denoted by $pf\delta\mathcal{P}cl(A)$ (resp. $pf\delta\mathcal{S}cl(A)$ and $pfecl(A)$) and are defined as follows: $pf\delta\mathcal{P}cl(A)$ (resp. $pf\delta\mathcal{S}cl(A)$ and $pfecl(A)$) $= \cap \{K | K \text{ is an } pf\delta\mathcal{P}cs \text{ (resp. } pf\delta\mathcal{S}cs, pfecs) \text{ and } A \subseteq K\}$.

Definition 2.11 [35] Let (X_1, Γ_P) be an $pfts$ and $A = \{ \langle a, \lambda_A(a), \mu_A(a) \rangle | a \in X_1 \}$ be an pfs in X_1 . A set A is said to be pf [(i)]

1. θ -interior of A (briefly, $pf\theta int(A)$) is defined by $pf\theta int(A) = \cup \{ pfint(B) : B \subseteq A \text{ \& } B \text{ is a } pfcs \text{ in } X_1 \}$.
2. θ -open set (briefly, $pf\theta os$) if $A = pf\theta int(A)$.
3. θ -semi open set (briefly, $pf\theta sos$) if $A \subseteq pfcl(pf\theta int(A))$.
4. M -open set (briefly, $pfMos$) if $A \subseteq pfcl(pf\theta int(A)) \cup pfint(pf\delta cl(A))$.

The complement of a $pfMos$ (resp. $pf\theta os$ & $pf\theta sos$) is called an pfM (resp. $pf\theta$ & $pf\theta\mathcal{S}$) closed set (briefly, $pfMcs$ (resp. $pf\theta cs$ & $pf\theta\mathcal{S}cs$)) in X_1 .

The family of all $pf\theta os$ (resp. $pf\theta cs, pf\theta sos, pf\theta\mathcal{S}cs, pfMos$ and $pfMcs$) of X_1 is denoted by $pf\theta OS(X_1)$, (resp. $pf\theta CS(X_1), pf\theta SOS(X_1), pf\theta\mathcal{S}CS(X_1), pfMOS(X_1)$ and $pfMCS(X_1)$).

Definition 2.12 [35] Let (X_1, Γ_P) be an $pfts$ and $A = \{ \langle a, \lambda_A(a), \mu_A(a) \rangle | a \in X_1 \}$ be an pfs in X_1 . Then the pf [(i)]

1. M (resp. $pf\theta$ -semi)-interior of A (briefly, $pfMint(A)$ (resp. $pf\theta Sint(A)$)) is defined by $pfMint(A)$ (resp. $pf\theta int(A)$ and $pf\theta Sint(A)$) $= \cup \{ B : B \subseteq A \text{ and } B \text{ is a } pfMos \text{ (resp. } pf\theta sos) \text{ in } X_1 \}$.
2. M (resp. θ -semi)-closure of A (briefly, $pfMcl(A)$ (resp. $pf\theta Scl(A)$)) is defined by $pfMcl(A)$ (resp. $pf\theta Scl(A)$) $= \cap \{ B : A \subseteq B \text{ and } A \text{ is a } pfMcs \text{ (resp. } pf\theta\mathcal{S}cs) \text{ in } X_1 \}$.

3 Pythagorean fuzzy contra M -open mappings

Definition 3.1 Let (X_1, Γ_P) and (X_2, Ψ_P) be any two $pfts$'s. A mapping $h_P: (X_1, \Gamma_P) \rightarrow (X_2, \Psi_P)$ is said to be a Pythagorean fuzzy (resp. δ , $\delta\mathcal{P}$, $\delta\mathcal{S}$, e , θ , $\theta\mathcal{S}$ and M)-continuous (briefly, $pfCts$ (resp. $pf\delta Cts$, $pf\delta\mathcal{P}Cts$, $pf\delta\mathcal{S}Cts$, $pfeCts$, $pf\theta Cts$, $pf\theta\mathcal{S}Cts$ and $pfMCts$)) if the inverse image of every $pfos$ in (X_2, Ψ_P) is a $pfos$ (resp. $pf\delta os$, $pf\delta\mathcal{P}os$, $pf\delta\mathcal{S}os$, $pfeos$, $pf\theta os$, $pf\theta\mathcal{S}os$ and $pfMos$) in (X_1, Γ_P) .

Definition 3.2 Let (X_1, Γ_P) and (X_2, Ψ_P) be any two $pfts$'s. A mapping $h_P: (X_1, \Gamma_P) \rightarrow (X_2, \Psi_P)$ is said to be a Pythagorean fuzzy (resp. θ , $\theta\mathcal{S}$, δ , $\delta\mathcal{P}$, $\delta\mathcal{S}$, M and e)-open (briefly, pfO (resp. $pf\theta O$, $pf\theta\mathcal{S}O$, $pf\delta O$, $pf\delta\mathcal{P}O$, $pf\delta\mathcal{S}O$, $pfMO$ and $pfeO$)) mapping if the image of every $pfos$ in (X_1, Γ_P) is a $pfos$ (resp. $pf\theta os$, $pf\theta\mathcal{S}os$, $pf\delta os$, $pf\delta\mathcal{P}os$, $pf\delta\mathcal{S}os$, $pfMos$ and $pfeos$) in (X_2, Ψ_P) .

Definition 3.3 Let (X_1, Γ_P) and (X_2, Ψ_P) be any two $pfts$'s. A mapping $h_P: (X_1, \Gamma_P) \rightarrow$

(X_2, Ψ_P) is said to be a Pythagorean fuzzy contra (resp. δ , $\delta\mathcal{P}$, $\delta\mathcal{S}$, e , θ , $\theta\mathcal{S}$ and M)-continuous (briefly, $pfcontraCts$ (resp. $pfcontra\delta Cts$, $pfcontra\delta\mathcal{P}Cts$, $pfcontra\delta\mathcal{S}Cts$, $pfcontraeCts$, $pfcontra\theta Cts$, $pfcontra\theta\mathcal{S}Cts$ and $pfcontraMCts$)) if the inverse image of every $pfos$ in (X_2, Ψ_P) is a $pfcs$ (resp. $pf\delta cs$, $pf\delta\mathcal{P}cs$, $pf\delta\mathcal{S}cs$, $pfecs$, $pf\theta cs$, $pf\theta\mathcal{S}cs$ and $pfMcs$) in (X_1, Γ_P) .

Definition 3.4 Let (X_1, Γ_P) and (X_2, Ψ_P) be any two pfts's. A mapping $h_P: (X_1, \Gamma_P) \rightarrow (X_2, \Psi_P)$ is said to be a Pythagorean fuzzy contra (resp. θ , $\theta\mathcal{S}$, δ , $\delta\mathcal{P}$, $\delta\mathcal{S}$, M and e)-open (briefly, $pfcontraO$ (resp. $pfcontra\theta O$, $pfcontra\theta\mathcal{S}O$, $pfcontra\delta O$, $pfcontra\delta\mathcal{P}O$, $pfcontra\delta\mathcal{S}O$, $pfcontraMO$ and $pfcontraeO$)) mapping if the image of every $pfos$ in (X_1, Γ_P) is a $pfcs$ (resp. $pf\theta cs$, $pf\theta\mathcal{S}cs$, $pf\delta cs$, $pf\delta\mathcal{P}cs$, $pf\delta\mathcal{S}cs$, $pfMcs$ and $pfecs$) in (X_2, Ψ_P) .

Proposition 3.1 Let (X_1, Γ_P) & (X_2, Ψ_P) be a pfts's. Let $h_P: (X_1, \Gamma_P) \rightarrow (X_2, \Psi_P)$ be a mapping. Then the following statements are hold for pfts, but not conversely.

1. Every $pfcontra\theta O$ is a $pfcontraO$.
2. Every $pfcontra\theta O$ is a $pfcontra\theta\mathcal{S}O$.
3. Every $pfcontra\theta\mathcal{S}O$ is a $pfcontraMO$.
4. Every $pfcontra\delta O$ is a $pfcontra\delta\mathcal{S}O$.
5. Every $pfcontra\delta O$ is a $pfcontra\delta\mathcal{P}O$.
6. Every $pfcontra\delta\mathcal{S}O$ is a $pfcontraeO$.
7. Every $pfcontra\delta\mathcal{P}O$ is a $pfcontraMO$.
8. Every $pfcontraMO$ is a $pfcontraeO$.
9. Every $pfcontra\delta O$ is a $pfcontraO$.

Proof.

1. Let B be a $pfos$ in (X_1, Γ_P) . Since h_P is $pfcontra\theta O$, $h_P(B)$ is $pf\theta cs$ in (X_2, Ψ_P) . Since every $pf\theta cs$ is a $pfcs$, $h_P(B)$ is a $pfcs$ in (X_2, Ψ_P) . Hence, h_P is a $pfcontraO$.

2. Let B be a $pfos$ in (X_1, Γ_P) . Since h_P is $pfcontra\theta O$, $h_P(B)$ is $pf\theta cs$ in (X_2, Ψ_P) . Since every $pf\theta cs$ is a $pf\theta\mathcal{S}cs$, $h_P(B)$ is a $pfcs$ in (X_2, Ψ_P) . Hence, h_P is a $pfcontra\theta\mathcal{S}O$.

3. Let B be a $pfos$ in (X_1, Γ_P) . Since h_P is $pfcontra\theta\mathcal{S}O$, $h_P(B)$ is $pf\theta\mathcal{S}cs$ in (X_2, Ψ_P) . Since every $pf\theta\mathcal{S}cs$ is a $pfMcs$, $h_P(B)$ is a $pfMcs$ in (X_2, Ψ_P) . Hence, h_P is a $pfcontraMO$.

4. Let B be a $pfos$ in (X_1, Γ_P) . Since h_P is $pfcontra\delta O$, $h_P(B)$ is $pf\delta cs$ in (X_2, Ψ_P) . Since every $pf\delta cs$ is a $pf\delta\mathcal{S}cs$, $h_P(B)$ is a $pf\delta\mathcal{S}cs$ in (X_2, Ψ_P) . Hence, h_P is a $pfcontra\delta\mathcal{S}O$.

5. Let B be a $pfos$ in (X_1, Γ_P) . Since h_P is $pfcontra\delta O$, $h_P(B)$ is $pf\delta cs$ in

(X_2, Ψ_P) . Since every $pf\delta cs$ is a $pf\delta\mathcal{P}cs$, $h_P(B)$ is a $pf\delta\mathcal{P}cs$ in (X_2, Ψ_P) . Hence, h_P is a $pfcontra\delta\mathcal{P}O$.


6. Let B be a $pfos$ in (X_1, Γ_P) . Since h_P is $pfcontra\delta SO$, $h_P(B)$ is $pf\delta\mathcal{S}cs$ in (X_2, Ψ_P) . Since every $pf\delta\mathcal{S}cs$ is a $pfecs$, $h_P(B)$ is a $pfecs$ in (X_2, Ψ_P) . Hence, h_P is a $pfcontraeO$.

7. Let B be a $pfos$ in (X_1, Γ_P) . Since h_P is $pfcontra\delta\mathcal{P}O$, $h_P(B)$ is $pf\delta\mathcal{P}cs$ in (X_2, Ψ_P) . Since every $pf\delta\mathcal{P}cs$ is a $pfMcs$, $h_P(B)$ is a $pfMcs$ in (X_2, Ψ_P) . Hence, h_P is a $pfcontraMO$.

8. Let B be a $pfos$ in (X_1, Γ_P) . Since h_P is $pfcontraMO$, $h_P(B)$ is $pfMcs$ in (X_2, Ψ_P) . Since every $pfMcs$ is a $pfecs$, $h_P(B)$ is a $pfecs$ in (X_2, Ψ_P) . Hence, h_P is a $pfcontraeO$.

9. Let B be a $pfos$ in (X_1, Γ_P) . Since h_P is $pfcontra\delta O$, $h_P(B)$ is $pf\delta cs$ in (X_2, Ψ_P) . Since every $pf\delta cs$ is a $pfcs$, $h_P(B)$ is a $pfcs$ in (X_2, Ψ_P) . Hence, h_P is a $pfcontraO$.

Remark 3.1 We obtain the following diagram from the results are discussed above.



Note: $A \rightarrow B$ denotes A implies B , but not conversely.

Example 3.1 Let $X_1 = X_2 = X = \{x_1, x_2\}$ and pfs 's A_1, A_2, A_3 & A_4 in X_1 , B_1, B_2, B_3 & B_4 in X_2 are defined as,

$$\begin{aligned} A_1 &= \{ \langle x_1, 0.20, 0.80 \rangle, \langle x_2, 0.40, 0.60 \rangle \} \\ A_2 &= \{ \langle x_1, 0.10, 0.90 \rangle, \langle x_2, 0.30, 0.70 \rangle \} \\ A_3 &= \{ \langle x_1, 0.90, 0.10 \rangle, \langle x_2, 0.70, 0.30 \rangle \} \\ A_4 &= \{ \langle x_1, 0.20, 0.80 \rangle, \langle x_2, 0.30, 0.70 \rangle \} \\ B_1 &= \{ \langle x_1, 0.80, 0.20 \rangle, \langle x_2, 0.60, 0.40 \rangle \} \end{aligned}$$

$$\begin{aligned} B_2 &= \{ \langle x_1, 0.90, 0.10 \rangle, \langle x_2, 0.70, 0.30 \rangle \} \\ B_3 &= \{ \langle x_1, 0.10, 0.90 \rangle, \langle x_2, 0.30, 0.70 \rangle \} \\ B_4 &= \{ \langle x_1, 0.80, 0.20 \rangle, \langle x_2, 0.70, 0.30 \rangle \} \end{aligned}$$

Now, we have $\Gamma_P = \{0_X, 1_X, A_1, A_2, A_3, A_4\}$ and $\Psi_P = \{0_X, 1_X, B_1, B_2, B_3, B_4\}$. Let $h_P: (X_1, \Gamma_P) \rightarrow (X_2, \Psi_P)$ be an identity mapping. Then, h_P is *pfcontraO* but not *pfcontraθO*, because the set A_1 is *pfos* in X_1 but $h_P(A_1) = A_1$ is not *pfθcs* in X_2 .

Example 3.2 Let $X_1 = X_2 = X = \{x_1, x_2\}$ and *pfs*'s A_1, A_2, A_3, A_4 in X_2 & B_1 in X_1 are defined as,

$$\begin{aligned} A_1 &= \{ \langle x_1, 0.80, 0.20 \rangle, \langle x_2, 0.60, 0.40 \rangle \} \\ A_2 &= \{ \langle x_1, 0.90, 0.10 \rangle, \langle x_2, 0.70, 0.30 \rangle \} \\ A_3 &= \{ \langle x_1, 0.10, 0.90 \rangle, \langle x_2, 0.30, 0.70 \rangle \} \\ A_4 &= \{ \langle x_1, 0.80, 0.20 \rangle, \langle x_2, 0.70, 0.30 \rangle \} \\ B_1 &= \{ \langle x_1, 0.80, 0.20 \rangle, \langle x_2, 0.60, 0.40 \rangle \} \end{aligned}$$

Now, we have $\Gamma_P = \{0_X, 1_X, B_1\}$ and $\Psi_P = \{0_X, 1_X, A_1, A_2, A_3, A_4\}$. Let $h_P: (X_1, \Gamma_P) \rightarrow (X_2, \Psi_P)$ be an identity mapping. Then, h_P is *pfcontraθSO* (resp. *pfcontraδSO*) but not *pfcontraθO* (resp. *pfcontraδO*), because the set B_1 is *pfos* in X_1 but $h_P(B_1) = B_1$ is not *pfθcs* (resp. *pfδcs*) in X_2 .

Example 3.3 Let $X_1 = X_2 = X = \{x_1, x_2\}$ and *pfs*'s A_1, A_2, A_3, A_4 in X_2 & B_1 in X_1 are defined as,

$$\begin{aligned} B_1 &= \{ \langle x_1, 0.20, 0.80 \rangle, \langle x_2, 0.40, 0.60 \rangle \} \\ A_1 &= \{ \langle x_1, 0.80, 0.20 \rangle, \langle x_2, 0.60, 0.40 \rangle \} \\ A_2 &= \{ \langle x_1, 0.90, 0.10 \rangle, \langle x_2, 0.70, 0.30 \rangle \} \\ A_3 &= \{ \langle x_1, 0.10, 0.90 \rangle, \langle x_2, 0.30, 0.70 \rangle \} \\ A_4 &= \{ \langle x_1, 0.80, 0.20 \rangle, \langle x_2, 0.70, 0.30 \rangle \} \end{aligned}$$

Now, we have $\Gamma_P = \{0_X, 1_X, B_1\}$ and $\Psi_P = \{0_X, 1_X, A_1, A_2, A_3, A_4\}$. Let $h_P: (X_1, \Gamma_P) \rightarrow (X_2, \Psi_P)$ be an identity mapping. Then, h_P is *pfcontraMO* but not *pfcontraθSO*, because the set B_1 is *pfos* in X_1 but $h_P(B_1) = B_1$ is not *pfθScs* in X_2 .

Example 3.4 Let $X_1 = X_2 = X = \{x_1, x_2\}$ and *pfs*'s A_1, A_2, A_3, A_4 in X_2 & B_1 in X_1 are defined as,

$$\begin{aligned} A_1 &= \{ \langle x_1, 0.80, 0.20 \rangle, \langle x_2, 0.60, 0.40 \rangle \} \\ A_2 &= \{ \langle x_1, 0.90, 0.10 \rangle, \langle x_2, 0.70, 0.30 \rangle \} \\ A_3 &= \{ \langle x_1, 0.10, 0.90 \rangle, \langle x_2, 0.30, 0.70 \rangle \} \\ A_4 &= \{ \langle x_1, 0.80, 0.20 \rangle, \langle x_2, 0.70, 0.30 \rangle \} \\ B_1 &= \{ \langle x_1, 0.40, 0.20 \rangle, \langle x_2, 0.40, 0.40 \rangle \} \end{aligned}$$

Now, we have $\Gamma_P = \{0_X, 1_X, B_1\}$ and $\Psi_P = \{0_X, 1_X, A_1, A_2, A_3, A_4\}$. Let $h_P: (X_1, \Gamma_P) \rightarrow (X_2, \Psi_P)$ be an identity mapping. Then, h_P is *pfcontraeO* but not *pfcontraMO*, because the set B_1 is *pfos* in X_1 but $h_P(B_1) = B_1$ is not *pfMcs* in X_2 .

Example 3.5 Let $X_1 = X_2 = X = \{x_1, x_2\}$ and pfs's A_1, A_2, A_3, A_4 in X_2 & B_1 in X_1 are defined as,

$$\begin{aligned} A_1 &= \{ \langle x_1, 0.80, 0.20 \rangle, \langle x_2, 0.60, 0.40 \rangle \} \\ A_2 &= \{ \langle x_1, 0.90, 0.10 \rangle, \langle x_2, 0.70, 0.30 \rangle \} \\ A_3 &= \{ \langle x_1, 0.10, 0.90 \rangle, \langle x_2, 0.30, 0.70 \rangle \} \\ A_4 &= \{ \langle x_1, 0.80, 0.20 \rangle, \langle x_2, 0.70, 0.30 \rangle \} \\ B_1 &= \{ \langle x_1, 0.20, 0.80 \rangle, \langle x_2, 0.30, 0.70 \rangle \} \end{aligned}$$

Now, we have $\Gamma_P = \{0_X, 1_X, B_1\}$ and $\Psi_P = \{0_X, 1_X, A_1, A_2, A_3, A_4\}$. Let $h_P: (X_1, \Gamma_P) \rightarrow (X_2, \Psi_P)$ be an identity mapping. Then, h_P is *pfcontra* \mathcal{O} (resp. *pfcontrae* \mathcal{O} and *pfcontra* $\delta\mathcal{PO}$) but not *pfcontra* $\delta\mathcal{O}$ (resp. *pfcontra* $\delta\mathcal{SO}$ and *pfcontra* $\delta\mathcal{O}$), because the set B_1 is *pfos* in X_1 but $h_P(B_1) = B_1$ is not *pf* δ *cs* (resp. *pf* δ *Scs* and *pf* δ *cs*) in X_2 .

Example 3.6 Let $X_1 = X_2 = X = \{x_1, x_2\}$ and pfs's A_1, A_2, A_3, A_4 in X_2 & B_1 in X_1 are defined as,

$$\begin{aligned} A_1 &= \{ \langle x_1, 0.80, 0.20 \rangle, \langle x_2, 0.60, 0.40 \rangle \} \\ A_2 &= \{ \langle x_1, 0.90, 0.10 \rangle, \langle x_2, 0.70, 0.30 \rangle \} \\ A_3 &= \{ \langle x_1, 0.10, 0.90 \rangle, \langle x_2, 0.30, 0.70 \rangle \} \\ A_4 &= \{ \langle x_1, 0.80, 0.20 \rangle, \langle x_2, 0.70, 0.30 \rangle \} \\ B_1 &= \{ \langle x_1, 0.80, 0.20 \rangle, \langle x_2, 0.60, 0.30 \rangle \} \end{aligned}$$

Now, we have $\Gamma_P = \{0_X, 1_X, B_1\}$ and $\Psi_P = \{0_X, 1_X, A_1, A_2, A_3, A_4\}$. Let $h_P: (X_1, \Gamma_P) \rightarrow (X_2, \Psi_P)$ be an identity mapping. Then, h_P is *pfcontra* \mathcal{MO} but not *pfcontra* $\delta\mathcal{PO}$, because the set B_1 is *pfcs* in X_1 but $h_P(B_1) = B_1$ is not *pf* $\delta\mathcal{Pcs}$ in X_2 .

Theorem 3.1 A mapping $h_P: (X_1, \Gamma_P) \rightarrow (X_2, \Psi_P)$ is *pfcontra* \mathcal{MO} iff for every pfs K of (X_1, Γ_P) , $h_P(pfint(K)) \supseteq pfMcl(h_P(K))$.

Proof. Necessity: Let h_P be a *pfMO* mapping and K be a *pfos* in (X_1, Γ_P) . Now, $pfint K \subseteq K$ implies $h_P(pfint(K)) \subseteq h_P(K)$. Since h_P is a *pfcontra* \mathcal{MO} mapping, $h_P(pfint(K))$ is *pfMcs* in (X_2, Ψ_P) such that $h_P(pfint(K)) \supseteq h_P(K)$. Therefore $h_P(pfint(K)) \supseteq pfMcl h_P(K)$.

Sufficiency: Assume K is a *pfos* of (X_1, Γ_P) . Then $h_P(K) = h_P(pfint(K)) \supseteq pfMcl h_P(K)$. But $pfMcl(h_P(K)) \supseteq h_P(K)$. So $h_P(K) = pfMcl(K)$ which implies $h_P(K)$ is a *pfMcs* of (X_2, Ψ_P) and hence h_P is a *pfcontra* \mathcal{MO} .

Theorem 3.2 If $h_P: (X_1, \Gamma_P) \rightarrow (X_2, \Psi_P)$ is a *pfcontra* \mathcal{MO} mapping, then $pfint(h_P^{-1}(K)) \subseteq h_P^{-1}(pfMcl(K))$ for every pfs K of (X_2, Ψ_P) .

Proof. Let K be a pfs of (X_2, Ψ_P) . Then $pfint(h_P^{-1}(K))$ is a *pfos* in (X_1, Γ_P) . Since h_P is *pfcontra* \mathcal{MO} , $h_P(pfint(h_P^{-1}(K)))$ is *pfMcs* in (X_2, Ψ_P) and hence $h_P(pfint(h_P^{-1}(K))) \subseteq pfMcl(h_P(h_P^{-1}(K))) \subseteq pfMcl(K)$. Thus $pfint(h_P^{-1}(K)) \subseteq h_P^{-1}(pfMcl(K))$.

Theorem 3.3 A mapping $h_P: (X_1, \Gamma_P) \rightarrow (X_2, \Psi_P)$ is *pfcontraMO* iff for each *pfs* G of (X_2, Ψ_P) and for each *pfos* K of (X_1, Γ_P) containing $h_P^{-1}(G)$, there is a *pfMos* H of (X_2, Ψ_P) such that $G \subseteq H$ and $h_P^{-1}(H) \subseteq K$.

Proof. Necessity: Assume h_P is a *pfcontraMO* mapping. Let G be the *pfcs* of (X_2, Ψ_P) and K is a *pfos* of (X_1, Γ_P) such that $h_P^{-1}(G) \subseteq K$. Then $H = (h_P(K^c))^c$ is *pfMos* of (X_2, Ψ_P) such that $h_P^{-1}(H) \subseteq K$.

Sufficiency: Assume K is a *pfos* of (X_1, Γ_P) . Then $h_P^{-1}((h_P(K))^c) \subseteq K^c$ and K^c is *pfcs* in (X_1, Γ_P) . By hypothesis, there is a *pfMos* H of (X_2, Ψ_P) such that $(h_P(K))^c \subseteq H$ and $h_P^{-1}(H) \subseteq K^c$. Therefore $K \subseteq (h_P^{-1}(H))^c$. Hence $H^c \subseteq h_P(K) \subseteq h_P((h_P^{-1}(H))^c) \subseteq H^c$ which implies $h_P(K) = H^c$. Since H^c is *pfMcs* of (X_2, Ψ_P) , $h_P(K)$ is *pfMc* in (X_2, Ψ_P) and thus h_P is *pfcontraMO* mapping.

Theorem 3.4 A mapping $h_P: (X_1, \Gamma_P) \rightarrow (X_2, \Psi_P)$ is *pfcontraMO* iff $h_P^{-1}(pfMcl(G)) \supseteq pfint(h_P^{-1}(G))$ for every *pfs* G of (X_2, Ψ_P) .

Proof. Necessity: Assume h_P is a *pfcontraMO* mapping. For any *pfs* G of (X_2, Ψ_P) , $h_P^{-1}(G) \subseteq pfcl(h_P^{-1}(G))$. Therefore by Theorem 3.3, there exists a *pfMos* K in (X_2, Ψ_P) such that $G \supseteq K$ and $h_P^{-1}(K) \supseteq pfint(h_P^{-1}(G))$. Therefore we obtain that $h_P^{-1}(pfMcl(G)) \supseteq h_P^{-1}(K) \supseteq pfint(h_P^{-1}(G))$.

Sufficiency: Assume G is a *pfs* of (X_2, Ψ_P) and K is a *pfcs* of (X_1, Γ_P) containing $h_P^{-1}(G)$. Put $H = pfcl(G)$, then $G \subseteq H$ and H is *pfMcs* and $h_P^{-1}(H) \subseteq pfint(h_P^{-1}(G)) \subseteq K$. Then by Theorem 3.3, h_P is *pfMO* mapping.

Theorem 3.5 If $h_P: (X_1, \Gamma_P) \rightarrow (X_2, \Psi_P)$ and $g_P: (X_2, \Psi_P) \rightarrow (X_3, \Phi_P)$ be two *pf* mappings and $g_P \circ h_P: (X_1, \Gamma_P) \rightarrow (X_3, \Phi_P)$ is *pfcontraMO*. If $g_P: (X_2, \Psi_P) \rightarrow (X_3, \Phi_P)$ is *pfcontraMIrr*, then $h_P: (X_1, \Gamma_P) \rightarrow (X_2, \Psi_P)$ is *pfMO* mapping.

Proof. Let K be a *pfos* in (X_1, Γ_P) . Then $(g_P \circ h_P)(K)$ is *pfMcs* of (X_3, Φ_P) because $g_P \circ h_P$ is *pfcontraMO* mapping. Since g_P is *pfcontraMIrr* and $(g_P \circ h_P)(K)$ is *pfMcs* of (X_3, Φ_P) , $g_P^{-1}((g_P \circ h_P)(K)) = h_P(K)$ is *pfMos* in (X_2, Ψ_P) . Hence h_P is *pfMO* mapping.

Theorem 3.6 If $h_P: (X_1, \Gamma_P) \rightarrow (X_2, \Psi_P)$ is *pfO* and $g_P: (X_2, \Psi_P) \rightarrow (X_3, \Phi_P)$ is *pfcontraMO* mappings, then $g_P \circ h_P: (X_1, \Gamma_P) \rightarrow (X_3, \Phi_P)$ is *pfcontraMO*.

Proof. Let K be a *pfos* in (X_1, Γ_P) . Then $h_P(K)$ is a *pfos* of (X_2, Ψ_P) because h_P is a *pfO* mapping. Since g_P is *pfcontraMO*, $g_P(h_P(K)) = (g_P \circ h_P)(K)$ is a *pfMcs* of (X_3, Φ_P) . Hence $g_P \circ h_P$ is *pfcontraMO* mapping.

4 Pythagorean fuzzy contra **M**-closed mapping

Definition 4.1 Let (X_1, Γ_P) and (X_2, Ψ_P) be any two *pfts*'s. A mapping $h_P: (X_1, \Gamma_P) \rightarrow (X_2, \Psi_P)$ is said to be a Pythagorean fuzzy (resp. θ , $\theta\mathcal{S}$, δ , $\delta\mathcal{P}$, $\delta\mathcal{S}$, **M** and **e**)-closed (briefly, *pfC* (resp. *pf θ C*, *pf $\theta\mathcal{S}$ C*, *pf δ C*, *pf $\delta\mathcal{P}$ C*, *pf $\delta\mathcal{S}$ C*, *pfMC* and *pfeC*)) mapping if the image of every *pfcs* in (X_1, Γ_P) is a *pfcs* (resp. *pf θ cs*, *pf $\theta\mathcal{S}$ cs*, *pf δ cs*, *pf $\delta\mathcal{P}$ cs*, *pf $\delta\mathcal{S}$ cs*, *pfMcs* and *pfecs*) in (X_2, Ψ_P) .

Definition 4.2 Let (X_1, Γ_P) and (X_2, Ψ_P) be any two pfts's. A mapping $h_P: (X_1, \Gamma_P) \rightarrow (X_2, \Psi_P)$ is said to be a Pythagorean fuzzy contra (resp. θ , $\theta\mathcal{S}$, δ , $\delta\mathcal{P}$, $\delta\mathcal{S}$, M and e)-closed (briefly, $pfcontraC$ (resp. $pfcontra\theta C$, $pfcontra\theta\mathcal{S}C$, $pfcontra\delta C$, $pfcontra\delta\mathcal{P}C$, $pfcontra\delta\mathcal{S}C$, $pfcontraMC$ and $pfcontraeC$)) mapping if the image of every pfcs in (X_1, Γ_P) is a pfos (resp. $pf\theta os$, $pf\theta\mathcal{S}os$, $pf\delta os$, $pf\delta\mathcal{P}os$, $pf\delta\mathcal{S}os$, $pfMos$ and $pfeos$) in (X_2, Ψ_P) .

Proposition 4.1 Let (X_1, Γ_P) & (X_2, Ψ_P) be a pfts's. Let $h_P: (X_1, \Gamma_P) \rightarrow (X_2, \Psi_P)$ be a mapping. Then the following statements are hold for pfts, but not conversely.

1. Every $pfcontra\theta C$ is a $pfcontraC$.
2. Every $pfcontra\theta C$ is a $pfcontra\theta\mathcal{S}C$.
3. Every $pfcontra\theta\mathcal{S}C$ is a $pfcontraMC$.
4. Every $pfcontra\delta C$ is a $pfcontra\delta\mathcal{S}C$.
5. Every $pfcontra\delta C$ is a $pfcontra\delta\mathcal{P}C$.
6. Every $pfcontra\delta\mathcal{S}C$ is a $pfcontraeC$.
7. Every $pfcontra\delta\mathcal{P}C$ is a $pfcontraMC$.
8. Every $pfcontraMC$ is a $pfcontraeC$.
9. Every $pfcontra\delta C$ is a $pfcontraC$.

Proof.

1. Let B be a pfcs in (X_1, Γ_P) . Since h_P is $pfcontra\theta C$, $h_P(B)$ is $pf\theta os$ in (X_2, Ψ_P) . Since every $pf\theta os$ is a $pfos$, $h_P(B)$ is a $pfos$ in (X_2, Ψ_P) . Hence, h_P is a $pfcontraC$.

2. Let B be a pfcs in (X_1, Γ_P) . Since h_P is $pfcontra\theta C$, $h_P(B)$ is $pf\theta os$ in (X_2, Ψ_P) . Since every $pf\theta os$ is a $pf\theta\mathcal{S}os$, $h_P(B)$ is a $pf\theta\mathcal{S}os$ in (X_2, Ψ_P) . Hence, h_P is a $pfcontra\theta\mathcal{S}C$.

3. Let B be a pfcs in (X_1, Γ_P) . Since h_P is $pfcontra\theta\mathcal{S}C$, $h_P(B)$ is $pf\theta\mathcal{S}os$ in (X_2, Ψ_P) . Since every $pf\theta\mathcal{S}os$ is a $pfMos$, $h_P(B)$ is a $pfMos$ in (X_2, Ψ_P) . Hence, h_P is a $pfcontraMC$.

4. Let B be a pfcs in (X_1, Γ_P) . Since h_P is $pfcontra\delta C$, $h_P(B)$ is $pf\delta os$ in (X_2, Ψ_P) . Since every $pf\delta os$ is a $pf\delta\mathcal{S}os$, $h_P(B)$ is a $pf\delta\mathcal{S}os$ in (X_2, Ψ_P) . Hence, h_P is a $pfcontra\delta\mathcal{S}C$.

5. Let B be a pfcs in (X_1, Γ_P) . Since h_P is $pfcontra\delta C$, $h_P(B)$ is $pf\delta os$ in (X_2, Ψ_P) . Since every $pf\delta os$ is a $pf\delta\mathcal{P}os$, $h_P(B)$ is a $pf\delta\mathcal{P}os$ in (X_2, Ψ_P) . Hence, h_P is a $pfcontra\delta\mathcal{P}C$.


6. Let B be a pfcs in (X_1, Γ_P) . Since h_P is $pfcontra\delta\mathcal{S}C$, $h_P(B)$ is $pf\delta\mathcal{S}os$ in (X_2, Ψ_P) . Since every $pf\delta\mathcal{S}os$ is a $pfeos$, $h_P(B)$ is a $pfeos$ in (X_2, Ψ_P) . Hence, h_P is a $pfcontraeC$.

7. Let B be a $pfcs$ in (X_1, Γ_P) . Since h_P is $pfcontra\delta PC$, $h_P(B)$ is $pf\delta Pos$ in (X_2, Ψ_P) . Since every $pf\delta Pos$ is a $pfMos$, $h_P(B)$ is a $pfMos$ in (X_2, Ψ_P) . Hence, h_P is a $pfcontraMC$.

8. Let B be a $pfcs$ in (X_1, Γ_P) . Since h_P is $pfcontraMC$, $h_P(B)$ is $pfMos$ in (X_2, Ψ_P) . Since every $pfMos$ is a $pfeos$, $h_P(B)$ is a $pfeos$ in (X_2, Ψ_P) . Hence, h_P is a $pfcontraeC$.

9. Let B be a $pfcs$ in (X_1, Γ_P) . Since h_P is $pfcontra\delta C$, $h_P(B)$ is $pf\delta os$ in (X_2, Ψ_P) . Since every $pf\delta os$ is a $pfos$, $h_P(B)$ is a $pfos$ in (X_2, Ψ_P) . Hence, h_P is a $pfcontraC$.

Remark 4.1 We obtain the following diagram from the results are discussed above.



Note: $A \rightarrow B$ denotes A implies B , but not conversely.

Example 4.1 Let $X_1 = X_2 = \{x_1, x_2\}$ and pfs 's A_1, A_2, A_3 & A_4 in X_1 , B_1, B_2, B_3 & B_4 in X_2 are defined as,

$$\begin{aligned} A_1 &= \{ \langle x_1, 0.20, 0.80 \rangle, \langle x_2, 0.40, 0.60 \rangle \} \\ A_2 &= \{ \langle x_1, 0.10, 0.90 \rangle, \langle x_2, 0.30, 0.70 \rangle \} \\ A_3 &= \{ \langle x_1, 0.90, 0.10 \rangle, \langle x_2, 0.70, 0.30 \rangle \} \\ A_4 &= \{ \langle x_1, 0.20, 0.80 \rangle, \langle x_2, 0.30, 0.70 \rangle \} \\ B_1 &= \{ \langle x_1, 0.80, 0.20 \rangle, \langle x_2, 0.60, 0.40 \rangle \} \\ B_2 &= \{ \langle x_1, 0.90, 0.10 \rangle, \langle x_2, 0.70, 0.30 \rangle \} \\ B_3 &= \{ \langle x_1, 0.10, 0.90 \rangle, \langle x_2, 0.30, 0.70 \rangle \} \\ B_4 &= \{ \langle x_1, 0.80, 0.20 \rangle, \langle x_2, 0.70, 0.30 \rangle \} \end{aligned}$$

Now, we have $\Gamma_P = \{0_X, 1_X, A_1, A_2, A_3, A_4\}$, $\Psi_P = \{0_X, 1_X, B_1, B_2, B_3, B_4\}$. Let $h_P: (X_1, \Gamma_P) \rightarrow (X_2, \Psi_P)$ be an identity mapping. Then, h_P is $pfcontraC$ but not $pfcontra\theta C$, because the set A_1^c is $pfcs$ in X_1 but $h_P(A_1^c) = A_1^c$ is not $pf\theta os$ in X_2 .

Example 4.2 Let $X_1 = X_2 = \{x_1, x_2\}$ and pfs 's A_1, A_2, A_3, A_4 in X_2 & B_1 in X_1 are defined as,

$$\begin{aligned} A_1 &= \{ \langle x_1, 0.80, 0.20 \rangle, \langle x_2, 0.60, 0.40 \rangle \} \\ A_2 &= \{ \langle x_1, 0.90, 0.10 \rangle, \langle x_2, 0.70, 0.30 \rangle \} \\ A_3 &= \{ \langle x_1, 0.10, 0.90 \rangle, \langle x_2, 0.30, 0.70 \rangle \} \\ A_4 &= \{ \langle x_1, 0.80, 0.20 \rangle, \langle x_2, 0.70, 0.30 \rangle \} \\ B_1 &= \{ \langle x_1, 0.80, 0.20 \rangle, \langle x_2, 0.60, 0.40 \rangle \} \end{aligned}$$

Now, we have $\Gamma_P = \{0_X, 1_X, B_1\}$ and $\Psi_P = \{0_X, 1_X, A_1, A_2, A_3, A_4\}$. Let $h_P: (X_1, \Gamma_P) \rightarrow (X_2, \Psi_P)$ be an identity mapping. Then, h_P is *pfcontra θ SC* (resp. *pfcontra δ SC*) but not *pfcontra θ C* (resp. *pfcontra δ C*), because the set B_1^c is *pfcs* in X_1 but $h_P(B_1^c) = B_1^c$ is not *pf θ os* (resp. *pf δ os*) in X_2 .

Example 4.3 Let $X_1 = X_2 = \{x_1, x_2\}$ and pfs 's A_1, A_2, A_3, A_4 in X_2 & B_1 in X_1 are defined as,

$$\begin{aligned} A_1 &= \{ \langle x_1, 0.80, 0.20 \rangle, \langle x_2, 0.60, 0.40 \rangle \} \\ A_2 &= \{ \langle x_1, 0.90, 0.10 \rangle, \langle x_2, 0.70, 0.30 \rangle \} \\ A_3 &= \{ \langle x_1, 0.10, 0.90 \rangle, \langle x_2, 0.30, 0.70 \rangle \} \\ A_4 &= \{ \langle x_1, 0.80, 0.20 \rangle, \langle x_2, 0.70, 0.30 \rangle \} \\ B_1 &= \{ \langle x_1, 0.20, 0.80 \rangle, \langle x_2, 0.40, 0.60 \rangle \} \end{aligned}$$

Now, we have $\Gamma_P = \{0_X, 1_X, B_1\}$ and $\Psi_P = \{0_X, 1_X, A_1, A_2, A_3, A_4\}$. Let $h_P: (X_1, \Gamma_P) \rightarrow (X_2, \Psi_P)$ be an identity mapping. Then, h_P is *pfcontraMC* but not *pfcontra θ SC*, because the set B_1^c is *pfcs* in X_1 but $h_P(B_1^c) = B_1^c$ is not *pf θ Sos* in X_2 .

Example 4.4 Let $X_1 = X_2 = \{x_1, x_2\}$ and pfs 's A_1, A_2, A_3, A_4 in X_2 & B_1 in X_1 are defined as,

$$\begin{aligned} A_1 &= \{ \langle x_1, 0.80, 0.20 \rangle, \langle x_2, 0.60, 0.40 \rangle \} \\ A_2 &= \{ \langle x_1, 0.90, 0.10 \rangle, \langle x_2, 0.70, 0.30 \rangle \} \\ A_3 &= \{ \langle x_1, 0.10, 0.90 \rangle, \langle x_2, 0.30, 0.70 \rangle \} \\ A_4 &= \{ \langle x_1, 0.80, 0.20 \rangle, \langle x_2, 0.70, 0.30 \rangle \} \\ B_1 &= \{ \langle x_1, 0.40, 0.20 \rangle, \langle x_2, 0.40, 0.40 \rangle \} \end{aligned}$$

Now, we have $\Gamma_P = \{0_X, 1_X, B_1\}$ and $\Psi_P = \{0_X, 1_X, A_1, A_2, A_3, A_4\}$. Let $h_P: (X_1, \Gamma_P) \rightarrow (X_2, \Psi_P)$ be an identity mapping. Then, h_P is *pfcontraeC* but not *pfcontraMC*, because the set B_1^c is *pfcs* in X_1 but $h_P(B_1^c) = B_1^c$ is not *pfMos* in X_2 .

Example 4.5 Let $X_1 = X_2 = \{x_1, x_2\}$ and pfs 's A_1, A_2, A_3, A_4 in X_2 & B_1 in X_1 are defined as,

$$\begin{aligned} A_1 &= \{ \langle x_1, 0.80, 0.20 \rangle, \langle x_2, 0.60, 0.40 \rangle \} \\ A_2 &= \{ \langle x_1, 0.90, 0.10 \rangle, \langle x_2, 0.70, 0.30 \rangle \} \end{aligned}$$

$$\begin{aligned} A_3 &= \{ \langle x_1, 0.10, 0.90 \rangle, \langle x_2, 0.30, 0.70 \rangle \} \\ A_4 &= \{ \langle x_1, 0.80, 0.20 \rangle, \langle x_2, 0.70, 0.30 \rangle \} \\ B_1 &= \{ \langle x_1, 0.20, 0.80 \rangle, \langle x_2, 0.30, 0.70 \rangle \} \end{aligned}$$

Now, we have $\Gamma_P = \{0_X, 1_X, B_1\}$ and $\Psi_P = \{0_X, 1_X, A_1, A_2, A_3, A_4\}$. Let $h_P: (X_1, \Gamma_P) \rightarrow (X_2, \Psi_P)$ be an identity mapping. Then, h_P is *pfcontraC* (resp. *pfcontraeC* and *pfcontra δ PC*) but not *pfcontra δ C* (resp. *pfcontra δ SC* and *pfcontra δ C*), because the set B_1 is *pfcs* in X_1 but $h_P(B_1^c) = B_1^c$ is not *pf δ os* (resp. *pf δ os* and *pf δ os*) in X_2 .

Example 4.6 Let $X_1 = X_2 = \{x_1, x_2\}$ and *pfs* 's A_1, A_2, A_3, A_4 in X_2 & B_1 in X_1 are defined as,

$$\begin{aligned} A_1 &= \{ \langle x_1, 0.80, 0.20 \rangle, \langle x_2, 0.60, 0.40 \rangle \} \\ A_2 &= \{ \langle x_1, 0.90, 0.10 \rangle, \langle x_2, 0.70, 0.30 \rangle \} \\ A_3 &= \{ \langle x_1, 0.10, 0.90 \rangle, \langle x_2, 0.30, 0.70 \rangle \} \\ A_4 &= \{ \langle x_1, 0.80, 0.20 \rangle, \langle x_2, 0.70, 0.30 \rangle \} \\ B_1 &= \{ \langle x_1, 0.80, 0.20 \rangle, \langle x_2, 0.60, 0.30 \rangle \} \end{aligned}$$

Now, we have $\Gamma_P = \{0_X, 1_X, B_1\}$ and $\Psi_P = \{0_X, 1_X, A_1, A_2, A_3, A_4\}$. Let $h_P: (X_1, \Gamma_P) \rightarrow (X_2, \Psi_P)$ be an identity mapping. Then, h_P is *pfcontraMC*, but not *pfcontra δ PC*, because the set B_1 is *pfcs* in X_1 but $h_P(B_1^c) = B_1^c$ is not *pf δ os* in X_2 .

Theorem 4.1 A mapping $h_P: (X_1, \Gamma_P) \rightarrow (X_2, \Psi_P)$ is *pfcontraMC* iff for each *pfs* G of (X_2, Ψ_P) and for each *pfos* K of (X_1, Γ_P) containing $h_P^{-1}(G)$, there is a *pfMcs* L of (X_2, Ψ_P) such that $G \subseteq L$ and $h_P^{-1}(L) \subseteq K$.

Proof. Necessity: Assume h_P is a *pfcontraMC* mapping. Let G be the *pfos* of (X_2, Ψ_P) and K is a *pfcs* of (X_1, Γ_P) such that $h_P^{-1}(G) \subseteq K$. Then $L = 1_Y - h_P^{-1}(K^c)$ is *pfMcs* of (X_2, Ψ_P) such that $h_P^{-1}(L) \subseteq K$.

Sufficiency: Assume K is a *pfcs* of (X_1, Γ_P) . Then $(h_P(K))^c$ is a *pfs* of (X_2, Ψ_P) and K^c is *pfos* in (X_1, Γ_P) such that $h_P^{-1}((h_P(K))^c) \subseteq K^c$. By hypothesis, there is a *pfMcs* L of (X_2, Ψ_P) such that $(h_P(K))^c \subseteq L$ and $h_P^{-1}(L) \subseteq K^c$. Therefore $K \subseteq (h_P^{-1}(L))^c$. Hence $L^c \subseteq h_P(K) \subseteq h_P((h_P^{-1}(L))^c) \subseteq L^c$ which implies $h_P(K) = L^c$. Since L^c is *pfMos* of (X_2, Ψ_P) , $h_P(K)$ is *pfMos* in (X_2, Ψ_P) and thus h_P is *pfcontraMC* mapping.

Theorem 4.2 If $h_P: (X_1, \Gamma_P) \rightarrow (X_2, \Psi_P)$ is *pfC* and $g_P: (X_2, \Psi_P) \rightarrow (X_3, \Phi_P)$ is *pfcontraMC*. Then $g_P \circ h_P: (X_1, \Gamma_P) \rightarrow (X_3, \Phi_P)$ is *pfcontraMC*.

Proof. Let K be a *pfcs* in (X_1, Γ_P) . Then $h_P(K)$ is *pfcs* of (X_2, Ψ_P) because h_P is *pfC* mapping. Now $(g_P \circ h_P)(K) = g_P(h_P(K))$ is *pfMos* in (X_3, Φ_P) because g_P is *pfcontraMC* mapping. Thus $g_P \circ h_P$ is *pfcontraMC* mapping.

Theorem 4.3 If $h_P: (X_1, \Gamma_P) \rightarrow (X_2, \Psi_P)$ is *pfcontraMC* map, then $pfMint(h_P(K)) \supseteq h_P(pfint(K))$.

Proof. Obvious.

Theorem 4.4 Let $h_P: (X_1, \Gamma_P) \rightarrow (X_2, \Psi_P)$ and $g_P: (X_2, \Psi_P) \rightarrow (X_3, \Phi_P)$ are *pfcontraMC* mappings. If every *pfMos* of (X_2, Ψ_P) is *pfos*, then $g_P \circ h_P: (X_1, \Gamma_P) \rightarrow$

(X_3, Φ_P) is *pfMC*.

Proof. Let K be a *pfcs* in (X_1, Γ_P) . Then $h_P(K)$ is *pfMos* of (X_2, Ψ_P) because h_P is *pfcontraMC* mapping. By hypothesis, $h_P(K)$ is *pfos* of (X_2, Ψ_P) . Now $g_P(h_P(K)) = (g_P \circ h_P)(K)$ is *pfMcs* in (X_3, Φ_P) because g_P is *pfcontraMC* mapping. Thus $g_P \circ h_P$ is *pfMC* mapping.

Theorem 4.5 Let $h_P: (X_1, \Gamma_P) \rightarrow (X_2, \Psi_P)$ be a bijective mapping. Then the following statements are equivalent: [(i)]

1. h_P is a *pfcontraMO* mapping.
2. h_P is a *pfcontraMC* mapping.
3. h_P^{-1} is *pfMCts* mapping.

Proof. (i) \Rightarrow (ii): Let us assume that h_P is a *pfcontraMO* mapping. By definition, K is a *pfos* in (X_1, Γ_P) , then $h_P(K)$ is a *pfMcs* in (X_2, Ψ_P) . Here, K is *pfcs* in (X_1, Γ_P) . Then $1_X - K$ is a *pfos* in (X_1, Γ_P) . By assumption, $h_P(1_X - K)$ is a *pfMcs* in (X_2, Ψ_P) . Hence, $1_Y - h_P(1_X - K)$ is a *pfMos* in (X_2, Ψ_P) . Therefore, h_P is a *pfcontraMC* mapping.

(ii) \Rightarrow (iii): Let K be a *pfcs* in (X_1, Γ_P) . By (ii), $h_P(K)$ is a *pfMos* in (X_2, Ψ_P) . Hence, $h_P(K) = (h_P^{-1})^{-1}(K)$. So h_P^{-1} is a *pfMos* in (X_2, Ψ_P) . Hence, h_P^{-1} is *pfMCts*.

(iii) \Rightarrow (i): Let K be a *pfos* in (X_1, Γ_P) . By (iii), $(h_P^{-1})^{-1}(K) = h_P(K)$ is a *pfcontraMO* mapping.

5 Conclusion

In this paper, the concepts of *pfcontraMO* and *pfcontraMC* mappings in *pfts* were discussed. Furthermore, the work was extended to include *pfcontraHom* and *pfcontraMHom*. In addition, the study demonstrated *pfcontraMCHom* and derived some of its related characteristics. In future, this work can be used in some mathematical applications.

References

- [1] K. T. Atanassov (1983), *ntuitionistic fuzzy sets*, VII ITKR Session, Sofia.
- [2] K. T. Atanassov (1986), *ntuitionistic fuzzy sets*, Fuzzy Sets Syst. **20**, 87-96.
- [3] K. T. Atanassov (1989), *Geometrical interpretation of the elements of the intuitionistic fuzzy objects*, Preprint IM-MFAIS-1-89, Sofia.
- [4] K. T. Atanassov (1999), *ntuitionistic fuzzy sets: theory and applications*, Physica, Heidelberg.
- [5] K. T. Atanassov (2012), *On intuitionistic fuzzy sets theory*, Springer, Berlin.
- [6] G. Beliakov and S. James (2014), *Averaging aggregation functions for preferences expressed as Pythagorean membership grades and fuzzy orthopairs*, In: Proceedings of the IEEE international conference on fuzzy systems (FUZZ-IEEE), 298-305.
- [7] B. Davvaz and E. H. Sadrabadi (2016), *An application of intuitionistic fuzzy sets in medicine*, Int. J. Biomath **9**, (3) 1650037.
- [8] S. K. De, R. Biswas and A. R. Roy (2001), *An application of intuitionistic fuzzy sets in medical diagnosis*, Fuzzy Sets Syst. **117** (2), 209-213.
- [9] S. Dick, R. R. Yager and O. Yazdanbakhsh (2016), *On Pythagorean and complex*

- fuzzy set operations. IEEE Trans Fuzzy Syst.* **24** (5), 1009-1021.
- [10] J. Dontchev, *contra-continuous functions and strongly S-closed spaces*, Internat. J. Maths. & Math. Sci., **19**, 303-310 (1996).
- [11] P. A. Ejegwa, A. J. Akubo and O. M. Joshua (2014), *intuitionistic fuzzy sets in career determination*, J Info Comput. Sci. **9** (4), 285-288.
- [12] P. A. Ejegwa (2015), *intuitionistic fuzzy sets approach in appointment of positions in an organization via max $\hat{=}$ min $\hat{=}$ max rule*, Glob. J Sci. Front Res F Math. Decis. Sci. **15** (6), 1-6.
- [13] P. A. Ejegwa and E. S. Modom (2015), *Diagnosis of viral hepatitis using new distance measure of intuitionistic fuzzy sets*, Int. J. Fuzzy Math. Arch. **8** (1), 1-7.
- [14] P. A. Ejegwa (2018), *Distance and similarity measures of Pythagorean fuzzy sets*, Granul Comput. <https://doi.org/10.1007/s41066-018-00149-z>.
- [15] H. Garg (2017), *A new improved score function of an interval-valued Pythagorean fuzzy set based TOPSnS method*, Int. J Uncertain Quantif **7** (5), 463-474.
- [16] H. Garg and S. Singh (2018), *A novel triangular interval type-2 intuitionistic fuzzy set and their aggregation operators*, Iran J Fuzzy Syst. **15** (5), 69-93.
- [17] H. Garg and K. Kumar (2018), *Distance measures for connection number sets based on set pair analysis and its applications to decision making process*, Appl. Intell **48** (10), 3346-3359.
- [18] H. Garg and K. Kumar (2018), *An advance study on the similarity measures of intuitionistic fuzzy sets based on the set pair analysis theory and their application in decision making*, Soft Comput. **22** (15), 4959-4970.
- [19] X. J. Gou, Z. S. Xu and P. J. Ren (2016), *The properties of continuous Pythagorean fuzzy information*, Int. J Intell. Syst. **31** (5), 401-424.
- [20] X. He, Y. Du and W. Liu (2016), *Pythagorean fuzzy power average operators*, Fuzzy Syst. Math. **30** (6), 116-124.
- [21] D. Jeeva, D. Sivakumar and A. Vadivel (2023), *Contra M -continuous Maps in Neutrosophic Soft Topological Spaces*, IJNS, **20** (3), 98-106.
- [22] D. Jeeva, A. Vadivel and D. Sivakumar (2023), *M -Continuous and M -nrresolute Maps in Neutrosophic Soft Topological Spaces*, AIP Conference Proceedings, 2852, 150001.
- [23] D. Jeeva, A. Vadivel and D. Sivakumar (2023), *Maps and homeomorphisms Via M -open sets in Neutrosophic Soft Topological Spaces*, South East Asian J. of Mathematics and Mathematical Sciences **19** (1) (2023), 367-384.
- [24] A. I. El-Maghrabi and M. A. Al-Juhani, *Mopen sets in topological spaces*, Pioneer J. Math. Sci., **4** (2) (2011), 213 $\hat{=}$ 230.
- [25] Murat Olgun, Mehmet Unver and Seyhmus Yardimci (2019), *Pythagorean fuzzy topological spaces*, Complex & Intelligent Systems. <https://doi.org/10.1007/s40747-019-0095-2>.
- [26] X. Peng and Y. Yang (2015), *Some results for Pythagorean fuzzy sets*, Int. J Intell Syst. **30**, 1133-1160.
- [27] X. Peng and G. Selvachandran (2017), *Pythagorean fuzzy set state of the art and future directions*, Artif Intell Rev. <https://doi.org/10.1007/s10462-017-9596-9>.
- [28] P. Revathi, K. Chitirakala and A. Vadivel, *Soft e -Separation Axioms in Neutrosophic soft Topological Spaces*, Journal of Physics: Conference Series, **2070**, (2021), 012028.
- [29] Supriti Saha, *Fuzzy δ -continuous mappings*, Journal of Mathematical Analysis and

- Applications, **126** (1987), 130-142.
- [30] E. Szmidt and J. Kacprzyk (2001), *intuitionistic fuzzy sets in some medical applications*, Note IFS **7** (4), 58-64.
 - [31] E. Szmidt and J. Kacprzyk (2004), *Medical diagnostic reasoning using a similarity measure for intuitionistic fuzzy sets*, Note IFS **10** (4), 61-69.
 - [32] M. Udhaya Shalini and A. Stanis Arul Mary (2022), *Generalized pre-closed sets in Pythagorean fuzzy topological spaces*, International Journal of Creative Research Thoughts (IJCRT), **10** (30), e142-e147.
 - [33] A. Vadivel, P. Manivannan, B. Vijayalakshmi and V. Chandrasekar, *Generalized Fuzzy contra e Continuous in Fuzzy Topological Spaces*, AIP Conference Proceedings, **2277**, 080001 (2020).
 - [34] A. Vadivel, M. Seenivasan and C. John Sundar, *An introduction to δ -open sets in a neutrosophic topological spaces*, Journal of Physics: Conference series, **1724** (2021), 012011.
 - [35] B. Vijayalakshmi, M. Ramalakshmi, A. Vadivel and G. Saravanakumar, *More on Maps and its Application in Pythagorean Fuzzy Topological Spaces*, accepted in Nanotechnology Perceptions.
 - [36] R. R. Yager (2013), *Pythagorean membership grades in multicriteria decision making*, In: Technical report MII -3301. Machine Intelligence Institute, Iona College, New Rochelle.
 - [37] R. R. Yager (2013), *Pythagorean fuzzy subsets*, In: Proceedings of the joint IFSA world congress NAFIPS annual meeting, 57-61.
 - [38] R. R. Yager (2014), *Pythagorean membership grades in multicriteria decision making*, IEEE Trans Fuzzy Syst. **22** (4), 958-965.
 - [39] L. A. Zadeh (1965), *Fuzzy sets*, Inf. Control, **8**, 338-353.