

# Experimental Methods For Evaluation Of Antidepressant Activity Using Laboratory Animal's Model

**Aswini Kumar Das, Pradeep Kumar Mohanty\***

*School of Pharmacy, LNCT University, Bhopal Madhya Pradesh India*

*Corresponding Author\**

*Pradeep Kumar Mohanty*

*Email: pmohanty1971@gmail.com*

*School of Pharmacy, LNCT University, Bhopal. (M.P.) India.*

Present investigation on the Experimental evaluation of hydroalcoholic extracts Chlorophytum borivilium roots (HCB) and n- butanol fractions of hydroalcoholic extract of Chorophytum borivilium roots (NHCB) were evaluated for antidepressant activity using in laboratory animals using different preclinical models of antidepressant activity such as forced swim test, tail suspension test and spontaneous locomotor activity. Both fractions showed potent antidepressant activity. The preliminary phytochemical analysis revealed the phytoconstituents present in the extracts. Extract of hydroalcoholic Chlorophytum borivilium roots (HCB) showed the presence of steroids, saponins, tannins, terpenoids, flavonoids, alkaloids. n- butanol fractions of hydroalcoholic extract of Chorophytum borivilium roots (NHCB) were subjected to the High-Performance Thin Layer Chromatography (HPTLC) studies hydroalcoholic extract of Chorophytum borivilium roots (NHCB) and their fractions showed potent antidepressant activity. The multistep putative action of these plants may be attributed to the prominent phytoconstituents namely 21-(4-hydroxy-2-methoxybenzofuran-5-yl)-3-phenylpropane-1, 3-dione from n-butanol fractions from n-butanol fractions of hydroalcoholic extracts of Chlorophytum borivilium roots (HCB). As a result of these studies two different phytoconstituents were isolated i.e., 1-(4-hydroxy-2-methoxybenzofuran-5-yl)-3-phenylpropane-1, 3-dione respectively from n-butanol fractions of hydroalcoholic extracts of Chlorophytum borivilium roots (HCB). The isolation of these phytoconstituents is an important concluding result of the study and antidepressant activity of the extracts may be attributed to these Phytoconstituents.

**Keywords:** Chlorophytum borivilium, CNS, Dysthymia, ANOVA, HPTLC.

## 1. INTRODUCTION

Depressive disorders are mood disorders that come in different forms, just as do other illnesses[1], such as heart disease and diabetes. Three of the most common types of depressive disorders are discussed below. Major depression is characterized by a combination of symptoms that last for at least two weeks in a row, including sad and/or irritable mood, that interfere with the ability to work, sleep, eat, and enjoy once pleasurable activities [2]. Difficulties in sleeping or eating can take the form of excessive or insufficient of either behaviour. Disabling episodes of depression can occur once, twice, or several times in a

lifetime. Depression affects not only the patients but also their friends and families. Nevertheless, depression can also occur in people without depression family history [3,4]. Recently diagnostic and statistical manual of mental disorder (DSMV) characterized major depression by symptoms: social withdrawal, lack of motivation, depressed mood, anhedonia, diminished interest in fun, sexual dysfunction, psychomotor agitation or retardation, fatigue, insomnia or hypersomnia, nightmares, guilt, inability to concentrate, sense of worthlessness and suicidal thinking. 15 to 25% of patients with depression have suicidal thoughts and sadism [5].

Today, a large variety of synthetic drugs are used as standard treatment for clinically depressed patients. Regardless of the medication that may be used to treat depression, practitioners have become more aware that different ethnic groups may have different responses and have different risks for side effects than others[6]. Antidepressant medications Selective serotonin reuptake inhibitors (SSRIs) are medications that increase the amount of the neurochemical serotonin in the brain. (Remember that brain serotonin levels are often low in depression.) As their name implies, the SSRIs work by selectively inhibiting (blocking) serotonin reuptake in the brain[7]. This block occurs at the synapse, the place where brain cells (neurons) are connected to each other. Serotonin is one of the chemicals in the brain that carries messages across these connections (synapses) from one neuron to another[8].

All the synthetic drugs available for the treatment of depression have various adverse effects and drug interactions. Such conditions create opportunities for alternative treatment of depression treatment[9]. Currently, researchers are seeking out more specific drugs with higher safety and lower cost. Traditional medicine offers several treatment options for mood disorders, most of them based on plant products which are empirically tested and proved to be safe in the past for human consumption[10]. Medicinal plants have attracted the attention of the researchers working in this field because these plants have long been used to treat different diseases, including psychiatric disorders and have less adverse effect than synthetic and chemical medicines. Several herbal drugs have been introduced during the recent past for decreasing depression in many emotional and physical disorders[11,12]. Medicinal herbs are still the preferred remedy for nearly 80% of people around the world, mainly in the developing countries to cure and improve the general health. This is primarily due to the common belief that plant derived drugs are without any adverse effects along with being economical and locally accessible[13]. In light of this, the present investigation is focused on the evaluation “Experimental methods for evaluation of Antidepressant activity using laboratory animal’s model”

## **2. MATERIALS AND METHODS**

### **Materials**

#### **Collection, identification and authentication of selected plant material**

Collection, identification and authentication of selected plant material *Chorophytum borivilium* roots were procured from Local area of Bhopal and were authenticated from Saifia

Science College, Bhopal MP, All the drugs and chemicals are AR grade and all the equipments/instruments used for the study were calibrated and validated[14].

### **Animals**

Swiss albino mice (20-40gm) and Wistar albino rats (180-220 gm) of either sex was procured from animal house as CCSEA reg 1587/PO/RE/S/11 and were maintained at  $25 \pm 2^\circ\text{C}$  and relative humidity of 45 to 55% and under standard environmental conditions (12 h light: 12 h dark cycle) at animal house of 100 . The animals had free access to food and water throughout study. Institutional Animal Ethical Committee approved the protocol. All the experiments were carried out between 9:00- 16:00 hour [15].

### **Methods:**

#### **Extraction of selected plant materials**

Powdered drugs (100 gm. each) were separately extracted successively using a Soxhlet extractor with hydroalcoholic solvent system is used in the ratio 1:1 and the temperature was set at 60-70 °C. Extracts were filtered, concentrated and after complete solvent evaporation, each of these solvent extracts was weighed and preserved at 5°C in an airtight bottle until further use[16]. Hydroalcoholic extract of *Chorophytum borivilium* roots was prepared and labeled as HCB.

#### **Preliminary phytochemical analysis of extracts [17]:**

The selected plant extracts were subjected to preliminary phytochemical analysis for the presence of prominent chemical constituents using following methods:

#### **Test for steroids**

##### **a. Salkowaski test**

2 ml of each Chloroform and concentrated sulphuric acid were added to this 2 ml of test solution, shaken and allowed to stand. Change in the colour of lower chloroform layer to red and acid layer to greenish yellow fluorescence which indicates the presence of steroids was observed[18].

##### **b. Liebermann-Burchard test**

2 ml test solution was mixed with chloroform (2 ml). To this solution, 2 ml of acetic anhydride and 2 drops of concentrated sulphuric acid from the side of test tube were added. Change in colour as, first red, then blue and finally green which indicates the presence of steroids was observed.

#### **• Test for triterpenoids**

##### **a. Salkowaski test**

2 ml of concentrated sulphuric acid was added to 2 ml of test solution. This mixture was shaken and allowed to stand. Change in the colour of lower layer to yellow which indicates the

presence of triterpenoids was observed.

#### **b. Liebermann-Burchardt Test**

The 3 ml of test solution was mixed with 3 ml of acetic anhydride, and then 2 ml of concentrated sulfuric acid was added to it from the sides of the test-tube. The development of deep red colour which indicates the presence of triterpenoids was observed.

#### • **Test for glycosides**

##### **a. Balget's test**

2 ml of the test solution was treated with 2 ml of sodium picrate solution. The development of yellow to orange colour which indicates the presence of cardiac glycosides was observed.

##### **b. Keller-Killiani test**

3-5 drops of glacial acetic acid, 1 drop of 5% FeCl<sub>3</sub>, and conc. sulphuric acid were added to the test tube containing 2 ml of test solution. Appearance of reddish-brown color at the junction of two layers and bluish green in the upper layer which indicates the presence of glycosides was observed.

##### **c. Legal's test**

To 2 ml of test solution, 1 ml of pyridine and 1 ml of sodium nitroprusside was added. Change in color to pink or red which indicates the presence of cardiac glycosides was observed.

##### **d. Borntrager's test**

2 ml of dilute sulphuric acid was added to 2 ml of test solution, boiled for a few minutes and filtered. To the filtrate 2 ml of benzene or chloroform was added and shaken well. The organic layer was separated and ammonia was added. The change in colour of ammoniacal layer to pink-red which indicates the presence of anthraquinone glycosides was observed.

#### • **Tests for saponins**

##### **a. Foam Test**

10 mg of extract was shaken vigorously with 1 ml water. Development of persistent foam which is stable at least for 15 minutes which indicates the presence of saponins was observed.

#### • **Tests for carbohydrates**

##### **a. Molisch's test**

3 ml of Molisch's reagent was added to the 3 ml of test solution, and was shaken for few minutes. Then 2 ml of concentrated sulphuric acid was added slowly from the sides of the test tube. The development of a purple ring at the junction of two liquids which indicates the presence of carbohydrates was observed.

##### **b. Barfoed's test**

1 ml of Barfoed's reagent and 1 ml of test solution were mixed in a test tube, heated in boiling

water bath for 2 minutes and then cooled. The appearance of red precipitate which indicates the presence of monosaccharides was observed.

**c. Fehling's test**

Fehling's A and B solutions (1 ml each) were added to the test tube and boiled for 1 minute. To this, 2 ml of test solution was added and heated in boiling water bath for 10 minutes. Appearance of yellow and then brick red precipitate which indicates the presence of reducing sugars was observed.

**d. Benedict's test**

Benedict's reagent (1 ml) and test solution (1ml) were mixed in a test tube and heated in boiling water bath for 10 minutes. Change in colour to yellow, green or red which indicates the presence of reducing sugar was observed.

• **Tests for alkaloids**

2 ml dilute hydrochloric acid was added to the 20 mg of dry extract, shaken well and filtered. With filtrate the following tests were performed.

**a. Mayer's test**

To the 3 ml of filtrate, 3 drops of Mayer's reagent (potassium mercuric iodide) was added. Appearance of reddish brown or cream precipitate which indicates the presence of alkaloids was observed.

**b. Hager's test**

To 3 ml of filtrate, 4-5 drops of Hager's reagent (saturated picric acid solution) was added. Appearance of yellow precipitate which indicates the presence of alkaloids was observed.

**c. Dragendorff's test**

3 ml of the filtrate was mixed with Dragendorff's reagent (potassium bismuth iodide). Appearance of reddish-brown precipitate which indicates the presence of alkaloids was observed.

• **Tests for Flavonoids**

**a. Ferric-chloride test**

Few drops (2-4) of ferric chloride solution were added to the test solution and appearance of intense green color which indicates the presence of flavonoids was observed.

**b. Shinoda test**

5 ml of ethanol (95%), 3 drops of hydrochloric acid and 0.5 gm magnesium turnings were added to 10 mg of extract. Change of color of solution to pink which indicates the presence of flavonoids was observed.

• **Tests for tannins**

**a. Ferric-chloride test**

3 ml of test solution was treated with 4-5 drops of ferric chloride solution. Development of dark color which indicates the presence of tannins was observed.

• **Tests for proteins**

**a. Millon's test**

Test solution (3 ml) and Million's reagent (5 ml) were mixed in a test tube. The appearance of white precipitate changing to brick red or dissolved and gave red color to solution on heating

which indicates the presence of proteins was observed.

**b. Xanthoproteic test**

To the test tube containing 3 ml test solution, 1 ml of conc. sulphuric acid was added. Appearance of white precipitate which turns yellow on boiling and orange on addition of  $\text{NH}_4\text{OH}$  which indicates the presence of tyrosin and/or tryptophan containing proteins was observed.

**d. Biuret test**

3 ml of the test solution was treated with 4% sodium hydroxide (3-5 drops) and 1% copper sulphate solution (3-5 drops). The appearance of blue colour which indicates the presence of proteins was observed.

**e. Ninhydrin test**

Test solution (3 ml) and 3 drops of 5% lead acetate solution were boiled on water bath for 10 min. Change in the color of solution to purple or blue which indicates the presence of amino acid was observed.

**Pharmacological screening of selected plant extracts[16]**

**Acute toxicity study of hydroalcoholic extracts of *Chlorophytum boriviliaum* roots (HCB) as per OECD guideline 425.**

Acute toxicity study was performed in healthy adult male albino mice (18-22 gm) as per guidelines (AOT 425) suggested by the Organization for Economical Co-operation and Development (OECD). Hydroalcoholic extract of ***Chlorophytum boriviliaum* roots (HCB)** were administered separately at the doses of 175, 550 and 2000 mg/kg in mice for oral toxicity study.

**Evaluation of antidepressant activity of hydroalcoholic extract of *Chlorophytum boriviliaum* roots (HCB) using Forced swim test**

30 Swiss albino mice of either sex are divided into eight groups and given the respective treatments for 14 days as follows.

Group-1: Animals of this group received distilled water (1 ml/kg, p.o.)

Group-2: Animals of this group received HCB (100 mg/kg, p.o.)

Group-3: Animals of this group received HCB (200 mg/kg, p.o.)

Group-4: Animals of this group received HCB (400 mg/kg, p.o.)

Group-5: Animals of this group received imipramine (5 mg/kg, p.o.)

Rats individually forced to swim in an open cylindrical container (diameter 10 cm, height 25 cm) containing 15 cm of water at  $25\pm 1^\circ\text{C}$ . All animals are forced to swim for 5 min and the duration of immobility is observed and measured. Each rat judged immobile when it ceased struggling and remained floating motionless in the water, making only those movements to keep its head above water[17]. A decrease in the duration of immobility is indicative of an antidepressant.

**Evaluation of antidepressant activity of hydroalcoholic extract of *Chlorophytum boriviliaum* roots (HCB) using Tail suspension test[18]**

48 Swiss albino mice of either sex are divided into eight groups and given the respective treatments for 14 days as follows.

Group-1: Animals of this group received distilled water (1 ml/kg, p.o.)

Group-2: Animals of this group received HCB (100 mg/kg, p.o.)

Group-3: Animals of this group received HCB (200 mg/kg, p.o.)

Group-4: Animals of this group received HCB (400 mg/kg, p.o.)

Group-5: Animals of this group received imipramine (5 mg/kg, p.o.)

On 14<sup>th</sup> day, one hour after the last dosing, mice were individually suspended on the edge of the table, 50 cm above the floor for the period of 10 minutes, with the help of adhesive tape placed approximately 1 cm from the tip of the tail. The total duration of immobility induced by tail suspension was recorded during a 6 min of the 10 min period. Mouse was considered immobile when it did not show any body movement, hanged passively and completely motionless.

### **Evaluation of antidepressant activity of hydroalcoholic extract of *Chlorophytum boriviliaum* roots (HCB) using spontaneous locomotor activity model[19]**

30 Swiss albino mice of either sex are divided into eight groups and given the respective treatments as follows.

Group-1: Animals of this group received distilled water (1 ml/kg, p.o.)

Group-2: Animals of this group received HCB (100 mg/kg, p.o.)

Group-3: Animals of this group received HCB (200 mg/kg, p.o.)

Group-4: Animals of this group received HCB (400 mg/kg, p.o.)

Group-5: Animals of this group received imipramine (5 mg/kg, p.o.)

On 14<sup>th</sup> day all the animals were subjected to assessment of locomotor activity using digital actophotometer. Then animals were placed in the digital actophotometer for 5 minutes and the locomotor activity counts displayed by interception of photobeams by movement of animals were be recorded and compare against control mice. Imipramine (5 mg/kg i.p.) was used as reference standard [34].

### **Evaluation of antidepressant activity of n- butanol fractions of n- butanol fractions of hydroalcoholic extract of *Chlorophytum boriviliaum* roots (NHCB) using Forced swim test**

24 Swiss albino mice of either sex are divided into six groups and given the respective treatments for 14 days as follows.

Group-1: Animals of this group received distilled water (1 ml/kg, p.o.)

Group-2: Animals of this group received NHCB (20 mg/kg, p.o.)

Group-3: Animals of this group received NHCB (40 mg/kg, p.o.)

Group-4: Animals of this group received imipramine (5 mg/kg, p.o.)

Rats individually forced to swim in an open cylindrical container (diameter 10 cm, height 25 cm) containing 15 cm of water at 25±1°C. All animals are forced to swim for 5 min and the duration of immobility is observed and measured. Each rat judged immobile when it ceased struggling and remained floating motionless in the water, making only those movements to keep its head above water. A decrease in the duration of immobility is indicative of an antidepressant.

### **Evaluation of antidepressant activity of n- butanol fractions of hydroalcoholic extract of *Chlorophytum borivilianum* roots (NHCB) using Tail suspension test**

24 Swiss albino mice of either sex are divided into six groups and given the respective treatments for 14 days as follows.

Group-1: Animals of this group received distilled water (1 ml/kg, p.o.)

Group-2: Animals of this group received NHCB (20 mg/kg, p.o.)

Group-3: Animals of this group received NHCB (40 mg/kg, p.o.)

Group-4: Animals of this group received imipramine (5 mg/kg, p.o.)

On 14<sup>th</sup> day, one hour after the last dosing, mice were individually suspended on the edge of the table, 50 cm above the floor for the period of 10 minutes, with the help of adhesive tape placed approximately 1 cm from the tip of the tail[20]. The total duration of immobility induced by tail suspension was recorded during a 6 min of the 10 min period. Mouse was considered immobile when it did not show any body movement, hanged passively and completely motionless.

### **Evaluation of antidepressant activity of n- butanol fractions of hydroalcoholic extract of *Chlorophytum borivilianum* roots (NHCB) using spontaneous locomotor activity model[21]**

36 Swiss albino mice of either sex are divided into six groups and given the respective treatments for 14 days as follows.

Group-1: Animals of this group received distilled water (1 ml/kg, p.o.)

Group-2: Animals of this group received NHCB (20 mg/kg, p.o.)

Group-3: Animals of this group received NHCB (40 mg/kg, p.o.)

Group-4: Animals of this group received imipramine (5 mg/kg, p.o.)

On 14<sup>th</sup> day all the animals were subjected to assessment of locomotor activity using digital actophotometer. Then animals were placed in the digital actophotometer for 5 minutes and the locomotor activity counts displayed by interception of photobeams by movement of animals were be recorded and compare against control mice. Imipramine (5 mg/kg i.p.) was used as reference standard [34].

### **Phytochemical qualitative analysis of extracts**

The selected plant extracts were subjected to phytochemical qualitative analysis for presence of various phytoconstituents of extracts.

### **High Performance Thin Layer Chromatography (HPTLC) studies of *Chlorophytum borivilianum* Fractions**

HPTLC system of CAMAG, Muttenz, Switzerland, Anchrom Enterprises (I) Pvt. Ltd, Mumbai, consisting of sample applicator (Linomat 5), Twin trough chamber with lid {10×10 cm, CAMAG, Muttenz, Switzerland), UV cabinet (Aetron, Mumbai) with dual wavelength (254/366 nm) and the HPTLC photodocumentation (Aetron, Mumbai) was used for study.

### **Chromatographic Conditions**

The samples were applied in the form of bands of width 6 mm with space between bands of 5 mm, with a 100 µL sample syringe (Hamilton, Bonaduz, Switzerland) on precoated silica gel aluminium plate 60 F254 (10 × 10, 5 × 10) with 250 µm thickness (E. MERCK, Darmstadt,



Germany) using a CAMAG Linomat 5 sample applicator (Switzerland). The plate was prewashed with methanol and activated at 110 °C for 5 minute, prior to chromatography[22]. The optimized chamber saturation time for mobile phase was kept 20 min. The length of chromatogram run was 8 cm. HPTLC plates were dried in a current of air with the help of a hair dryer. The slit dimensions of 5×0.45 mm and scanning speed of 20 mm/sec was employed in analysis.

### 3. RESULTS AND DISCUSS

#### Preliminary phytochemical analysis of extracts

**Table 1: Preliminary phytochemical evaluation of extracts**

Plant constituents	Tests performed	HCB
Test for Steroids	1. Salkowaski Test	++
	2. Liebermann-Buchard Test	++
Test for Triterpenoids	1. Salkowaski Test	++
	2. Liebermann-Buchard Test	++
Test for Glycosides	1. Balget's test	-
	2. Keller-Killiani test	-
	3. Legals test	-
	4. Borntrager's test	-
Tests for Saponins	1. Foam Test	++
Tests for Carbohydrates	1. Molisch's test	++
	2. Barfoed's test	++
	3. Fehling's test	++
	4. Benedict's test	++
Test for Alkaloids	1. Mayer's Reagent	++
	2. Hager's Reagent	++
	3. Dragendorff's Reagent	++
Tests for Flavonoids	1. Ferric-chloride test	++
	2. Shinoda test	++
Test for Tannins	1. FeCl <sub>3</sub> Solution	++
	2. Gelatin test	++
Test for Proteins	1. Millon's test	++
	2. Xanthoproteic test	++
	3. Biuret test	++
	4. Ninhydrin test	++

+ Present, - Absent (+ Small, ++Moderate, +++High)

The preliminary phytochemical analysis revealed the phytoconstituents present in the extracts. Hydroalcoholic extract of fruits of *Chlorophytum borivilium* roots (HCB) showed the presence of alkaloids, flavonoids, tannins, resins, carbohydrates, proteins and saponins.

### Pharmacological screening of selected plant extracts

#### Acute toxicity study of hydroalcoholic extract of *Chlorophytum borivilium* roots (HCB) as per OECD guideline 425.

There were no signs of morbidity nor mortality and any type of adverse effects found up to the dose of 2000 mg/kg p.o. for both the extracts[23].

**Table 2: Evaluation of antidepressant activity of hydroalcoholic extract of *Chlorophytum borivilium* roots (HCB) using Forced swim test**

S.No.	Treatment	Immobility time (secs) Mean±SEM
1	Normal control	75.78±2.63
2	HCB (100 mg/kg)	72.11±2.48
3	HCB (200 mg/kg)	70.09±3.08
4	HCB (400 mg/kg)	64.31±1.85**
5	Imipramine (5 mg/kg, p.o.)	53.48±2.25**

Values are mean ± S.E.M., n=6; Statistical analysis by one-way ANOVA followed by Dunnett's test using Graph pad Instat software; p<0.05\*, p<0.01\*\* and p<0.001\*\*\* compared to control group.

In this study, HCB 200 and 400 mg/kg showed dose dependent decrease in the immobility time as compared to normal control. Higher doses of both the extracts were found to be equipotent to that of reference standard i.e. imipramine[24].

**Table 3: Evaluation of antidepressant activity of hydroalcoholic extract of *Chlorophytum borivilium* roots (HCB) using Tail suspension test**

S. No.	Treatment	Immobility time (secs) Mean±SEM
1	Normal control	72.04±1.73
2	HCB (100 mg/kg)	67.71±1.84*
3	HCB (200 mg/kg)	62.96±2.49**
4	HCB (400 mg/kg)	57.08±2.19**
5	Imipramine (5 mg/kg, p.o.)	51.64±2.73**

Values are mean ± S.E.M., n=6; Statistical analysis by one-way ANOVA followed by Dunnett's test using Graph pad Instat software; p<0.05\*, p<0.01\*\* and p<0.001\*\*\* compared to control group.

In this study, all the doses of the extracts have shown significant decrease in the immobility time.

**Table 4: Evaluation of antidepressant activity of hydroalcoholic extract of Chlorophytum borivilium roots (HCB) using spontaneous locomotor activity model**

S. No.	Treatment	Locomotor score Mean±SEM
1	Normal control	42.00±2.73
2	HCB (100 mg/kg)	41.00±2.80
3	HCB (200 mg/kg)	42.00±03.37
4	HCB (400 mg/kg)	48.66±1.56
5	Imipramine (5 mg/kg, p.o.)	32.33±2.06

Values are mean ± S.E.M., n=6; Statistical analysis by one-way ANOVA followed by Dunnett's test using Graph pad InStat software; p<0.05\*, p<0.01\*\*and p<0.001\*\*\* compared to control group.

In this study, no treatment including Reference standard has shown any significant change in the locomotor activity. However, reference std has shown trend of reducing in the score.

**Table 5: Evaluation of antidepressant activity of n- butanol fractions of hydroalcoholic extract of Chlorophytum borivilium roots (NHCB) using Forced swim test**

S. No.	Treatment	Immobility time (secs) Mean±SEM
1	Normal control	64.16±1.63
2	NHCB (20 mg/kg)	46.02±1.29**
3	NHCB (40 mg/kg)	45.16±1.57**
4	Imipramine (5 mg/kg, p.o.)	42.68±0.80**

Values are mean ± S.E.M., n=6; Statistical analysis by one-way ANOVA followed by Dunnett's test using Graph pad InStat software; p<0.05\*, p<0.01\*\*and p<0.001\*\*\* compared to control group.

In this study, NHCB-20 and NHCB-40 showed significant decrease in the immobility time as compared to normal control. Furthermore, all these doses were equipotent to reference standard Imipramine.

**Table 6: Evaluation of antidepressant activity of n- butanol fractions of hydroalcoholic extract of Chlorophytum borivilium roots (NHCB) using Tail suspension test**

S. No.	Treatment	Immobility time (secs) Mean±SEM
1	Normal control	77.88±2.79
2	NETT (20 mg/kg)	70.98±1.98*
3	NETT (40 mg/kg)	67.95±1.17**
4	NHCB (20 mg/kg)	71.12±1.83**

5	NHCB (40 mg/kg)	64.30±1.56**
6	Imipramine (5 mg/kg, p.o.)	63.11±1.77**

Values are mean ± S.E.M., n=6; Statistical analysis by one-way ANOVA followed by Dunnett's test using Graph pad Instat software; p<0.05\*, p<0.01\*\* and p<0.001\*\*\* compared to control group.

In this study, extracts have shown significant dose dependent reduction in immobility time. The reference standard has shown significant action, which was statistically equal as that of higher dose of both fractions.

**Table 7: Evaluation of antidepressant activity of n- butanol fractions of hydroalcoholic extract of Chlorophytum boriviliaum roots (NHCB) using spontaneous locomotor activity model**


S. No.	Treatment	Locomotor score Mean±SEM
1	Normal control	49.66±1.47
2	NHCB (20 mg/kg)	50.00±1.46
3	NHCB (40 mg/kg)	49.83±1.30
4	Imipramine (5 mg/kg, p.o.)	43.66±1.43*

Values are mean ± S.E.M., n=6; Statistical analysis by one-way ANOVA followed by Dunnett's test using Graph pad Instat software; p<0.05\*, p<0.01\*\* and p<0.001\*\*\* compared to control group.


In this study, no treatment including Reference standard has shown any significant change in the locomotor activity. However, reference standard Imipramine has shown significant reduction.

#### **Phytochemical qualitative analysis of extracts**



The selected plant extracts were subjected to phytochemical qualitative analysis for presence of various phytoconstituents of extracts.



**Figure 1: High Performance Thin Layer Chromatography (HPTLC) studies of n- butanol fractions of hydroalcoholic extract of *Chlorophytum borivilium* roots**



B



**Figure 2:** n-Butanol IIb fraction of Hydroalcoholic extract of *Chlorophytum borivillanum* A) at 366 nm and B) at 254 nm and C) Visible light, Volume applied 10  $\mu$ l

and 20  $\mu$ l





Figure 3: FT-IR Spectrum of compound in n-butanol fraction Band 3 (Rf. 0.82)

S. No.	Part of molecule	Vibration	Frequency $\text{cm}^{-1}$
1	Aliphatic $\text{CH}_2 / \text{CH}_3$	a) C-H str	2991
		b) C-H bend	1446
2	-C(O)- $\text{CH}_2$ -	a) C=O str	1715
			1031,1262
3	-O- $\text{CH}_3$	a) C-O str	1031,1262
4	-C=C- ring (Furan ring)	a) C=C str	1623
		b) C-O str	1372
5	Ar ring	a) C-H str	3151
		b) C=C str	1524
		c) C-H bend	723
6	-OH	a) O-H str	3283



**Figure 4:** Mass Spectrum of compound in n-butanol fraction Band 3 (Rf. 0.82)


**Table 8:** IR data interpretation Band 3 (Rf. 0.82)

S. No.	$\delta$	Protons	Type
1	3.688	3 (s)	Methyl protons of $-O-CH_3$
2	4.041	2 (s)	Methylene protons $-CH_2-$
3	5.281	1 (s)	Proton on Furan ring of fused Benzofuran ring
4	6.809-6.971	1 (d)	Proton on Benzene ring of fused Benzofuran ring (a)
5	6.977-7.144	1 (d)	Proton on Benzene ring of fused Benzofuran ring (b)
6	7.212-7.857	5 (m)	Protons of phenyl ring
7	10.103	1 (s)	OH proton


**Table 9:** NMR data of compound in n-butanol fraction band 3 (Rf. 0.82)

**Conclusion:** Probable structure of the compound from IR, Mass and NMR is Probable structure of the compound from elemental analysis, IR and NMR is





**Figure 6: 1-(4-hydroxy-2-methoxybenzofuran-5-yl)-3-phenylpropane-1,3-dione**



**Figure 7: High Performance Thin Layer Chromatography (HPTLC) studies of n-Butanol IIb fraction of Hydroalcoholic extract of Chlorophytum borivillanum**

### Discussion

Depression disorder is a type of depression with a high degree of severity and accompanying symptoms and is one of the most common psychiatric disorders, with a prevalence of 2–20% in the general population.<sup>1</sup> According to statistics, depression ranks as the sixth leading cause of disease in the 10–49 years age group, affecting about 17% of the population[25], with a considerable disease burden.<sup>4</sup> Suicide rates are generally elevated in MDD, especially among younger patients, with at least 10% experiencing suicidal ideation and suicidal behavior.

Antidepressant medication (ADM) is currently one of the most commonly prescribed medications for MDD, with 10% or more of the general population taking antidepressants yearly in some high-income countries[26]. However, side effects of antidepressants are

frequently reported in studies, including central nervous system abnormalities, gastrointestinal reactions, weight changes, and allergic reactions.

Recently, herbal medicines have received much attention as alternative treatments for depression because of their fewer side effects and lower costs[27]. The description of the antidepressant activity of selected plant-derived compounds involves several mechanisms, including inhibition of monoamine reuptake; enhanced serotonin receptor binding and sensitization; monoamine oxidase inhibition; GABAergic effects (especially for plants exhibiting sedative and anxiolytic effects accompanying the antidepressant effect); complex, excitatory or inhibitory effects on various receptors (N-methyl-D-aspartic acid (NMDA), GABA, cholinergic, adrenergic, serotonergic, dopaminergic and opioid ones); and cannabinoid system effects[28,29].

Chlorophytum borivilianum are herbal medicines with wide range of pharmacological actions. These plants have been traditionally claimed to be effective in relieving depressed mood and anxiety and other such depressive disorders but have not been scientifically studied for their antidepressant activity[30].

The most trustworthy way to determine whether phytoconstituents are present is by a preliminary phytochemical examination, followed by a comparison of the results with existing literature[31]. So as to assess the phytochemical profile of the plants, hydroalcoholic extract of Chlorophytum borivilianum roots were subjected to preliminary phytochemical analysis. The preliminary phytochemical analysis revealed the phytoconstituents present in the extracts. Hydroalcoholic extract of fruits of Chlorophytum borivilianum roots (HCB) showed the presence of alkaloids, flavonoids, tannins, resins, carbohydrates, proteins and saponins[32]. These phytoconstituents have been reported to possess antidepressant effects.

To confirm the extent of a therapeutic drug's therapeutic use, toxicity profile testing is crucial for all therapeutic drugs[33]. Moreover, in the present investigation, the results of preliminary phytochemical analysis suggested further pharmacological exploration of extracts, hence it was essential to assess the extracts for their toxicity profile and thereby confirmation of their safety (OECD 2008). Hence the acute toxicity study of ethanolic extract of hydroalcoholic extract of Chlorophytum borivilianum roots (HCB) was carried out as per OECD guideline 425. The results of acute oral toxicity studies revealed that these extracts are safe with no signs of morbidity nor mortality and any type of adverse effects up to 2000 mg/kg which is highest prescribed limit as per this test[34]. Thus, the extracts fulfilled the safety criteria before the assessment of preclinical activity (OECD guidelines 425). Based upon these findings and available literature, the three different doses i.e. 100, 200 and 400 mg/kg of each extract were selected for the further preclinical investigations.

In the present study, the antidepressant effects of the the extracts were evaluated using three widely used tests i.e. forced swim test, tail suspension test and spontaneous locomotor activity test. These models are known to be of highly predictive relevance with respect to the clinical spectrum of activity[35].

In forced swim test, HCB 200 and 400 mg/kg showed dose dependent decrease in the immobility time as compared to normal control. The HCB 400 mg/kg is also found to be effective to reduce the immobility. Higher doses of the extracts were found to be equipotent to that of reference standard i.e. imipramine.

In the tail suspension test, all the doses of the extracts have shown significant decrease in the

immobility[36].

In spontaneous locomotor activity, no treatment including reference standard showed any significant change in the locomotor activity. However, reference standard Imipramine showed a non-significant reduction in locomotor activity.

After the pharmacological activity of individual extracts, the fractionation of the extracts was carried out and n-butanol fractions of hydroalcoholic extract of *Chorophytum borivilium* roots (NHCB) were prepared. In order to assess the antidepressant activity of these fractions, they were also subjected to the same preclinical models of antidepressant activity at the doses of 20 mg/kg and 40 mg/kg.

In the forced swim test, NHCB 20 mg/kg and NHCB 40 mg/kg showed significant decrease in the immobility time as compared to normal control. Furthermore, all these doses were equipotent to reference standard Imipramine.

In the tail suspension test, the fractions showed significant and dose dependent reduction in immobility time at both the doses. The fractions were found to be almost equipotent to each other. The reference standard has shown significant action, which was statistically equal as that of higher dose of fractions.

In the spontaneous locomotor activity, none of the fractions showed any significant change in the locomotor activity. However, reference standard Imipramine showed significant reduction in locomotor activity. Utilizing medicinal biodiversity to its fullest extent in order to search for phytochemicals with antidepressant properties is a difficult task. Additionally, administration of phytoconstituents without desired pharmacological activity led to precipitation of adverse effects or reduced therapeutic efficacy.

This problem can be overcome by the isolation of most active Phytoconstituents responsible for the desired pharmacological activity from the extract. This not only identifies and isolates the active Phytoconstituents but also provides an important input to the synthetic chemistry regarding the structure of patient friendly bioactive constituent with required pharmacological action. HPTLC technique n-butanol fractions of ethanolic extract of n-butanol fractions of hydroalcoholic extract of *Chorophytum borivilium* roots (NHCB) were subjected to the High-Performance Thin Layer Chromatography (HPTLC) studies.

As a result of these studies two different phytoconstituents were isolated i.e., 1-(4-hydroxy-2-methoxybenzofuran-5-yl)-3-phenylpropane-1, 3-dione from n-butanol fractions of hydroalcoholic extract of *Chorophytum borivilium* roots (NHCB). The isolation of these phytoconstituents is an important concluding result of the study and cardioprotective activity of the two extracts may be attributed to these phytoconstituents.

#### **4. Conclusion**

The present study the antidepressant activity of hydroalcoholic extract of *Chlorophytum borivilianum* roots and n-butanol fractions of hydroalcoholic extract of *Chorophytum borivilium* roots (NHCB) were evaluated for antidepressant activity using in laboratory animals using different preclinical models of antidepressant activity such as forced swim test, tail suspension test and spontaneous locomotor activity. Both fractions showed potent antidepressant activity. The multistep putative action of these plants may be attributed to the prominent phytoconstituents namely 1-(4-hydroxy-2-methoxybenzofuran-5-yl)-3-phenylpropane-1,3-dione respectively.

## Reference

1. Zhang, Z.; Deng, T.; Wu, M.; Zhu, A.; Zhu, G. Botanicals as modulators of depression and mechanisms involved. *Chin. Med.* 2019, 14, 24.
2. Monroe, SM, and Harkness, KL. Major depression and its recurrences: Life course matters. *Annu Rev Clin Psychol.* (2022) 18:329–57.
3. Ștefănescu R, Tero-Vescan A, Negroiu A, Aurică E, Vari C-E. A Comprehensive Review of the Phytochemical, Pharmacological, and Toxicological Properties of *Tribulus terrestris* L. *Biomolecules.* 2020; 10(5):752.
4. Khanam Z, Singh O, Singh R, Bhat IU. Safed musli (*Chlorophytum borivilianum*): a review of its botany, ethnopharmacology and phytochemistry. *J Ethnopharmacol.* 2013 Nov 25;150(2):421-41.
5. Sarris, J.; Panossian, A.; Schweitzer, I.; Stough, C.; Scholey, A. Herbal medicine for depression, anxiety and insomnia: A review of psychopharmacology and clinical evidence. *Eur. Neuropsychopharmacol.* 2011, 21, 841–860.
6. Wang, Y.S.; Shen, C.Y.; Jiang, J.G. Antidepressant active ingredients from herbs and nutraceuticals used in TCM: Pharmacological mechanisms and prospects for drug discovery. *Pharmacol. Res.* 2019, 150, 104520.
7. Farahani, M.S.; Bahramsoltani, R.; Farzaei, M.H.; Abdollahi, M.; Rahimi, R. Plant-derived natural medicines for the management of depression: An overview of mechanisms of action. *Rev. Neurosci.* 2015, 26, 305–321.
8. Lundstrom, K.; Pham, H.T.; Dinh, L.D. Interaction of plant extracts with central nervous system receptors. *Medicines* 2017, 4, 12
9. Zhu W, Du Y, Meng H, Dong Y, Li L. A review of traditional pharmacological uses, phytochemistry, and pharmacological activities of *Tribulus terrestris*. *Chem Cent J.* 2017 Jul 11;11(1):60.
10. Ștefănescu R, Tero-Vescan A, Negroiu A, Aurică E, Vari C-E. A Comprehensive Review of the Phytochemical, Pharmacological, and Toxicological Properties of *Tribulus terrestris* L. *Biomolecules.* 2020; 10(5):752.
11. Khanam Z, Singh O, Singh R, Bhat IU. Safed musli (*Chlorophytum borivilianum*): a review of its botany, ethnopharmacology and phytochemistry. *J Ethnopharmacol.* 2013 Nov 25;150(2):421-41.
12. Thakur GS, Bag M, Sanodiya BS, Debnath M, Zacharia A, Bhadauriya P, Prasad GB, Bisen PS. *Chlorophytum borivilianum*: a white gold for biopharmaceuticals and nutraceuticals. *Curr Pharm Biotechnol.* 2009 Nov;10(7):650-66.
13. OECD (2008). "Test No. 425: Acute Oral Toxicity: UpandDown Procedure. Paris, France: OECD Publishing."
14. Gambarana C, Scheggi S, Tagliamonte A, Tolu P, De Montis MG. Animal models for the study of antidepressant activity. *Brain Res Brain Res Protoc.* 2001 Apr;7(1):11-20.
15. Krishnan V, Nestler EJ. Animal models of depression: molecular perspectives. *Curr Top Behav Neurosci.* 2011;7:121-47.
16. Hildebert Wagner, Sabine Bladt, plant drug analysis, A thin layer chromatography Atlas Springer, Paris, 2nd edition, Page no:125, 1995.
17. Martins, J.; Brijesh, S. Phytochemistry and pharmacology of anti-depressant medicinal plants: A review. *Biomed. Pharmacother.* 2018, 104, 343–365.
18. Staines, S.S. Herbal medicines: Adverse effects and drug-herbs interactions. *J. Malta Coll. Pharm. Pract.* 2011, 17, 38–42
19. Sethi P.D. first ed. CBS Publishers and Distributors; New Delhi: 1996. High Performance Thin Layer Chromatography; pp. 1–56.
20. Gupta M, Thakur S, Sharma A, Gupta S. Qualitative and Quantitative Analysis of Phytochemicals and Pharmacological Value of Some Dye Yielding Medicinal Plants. *Orient J Chem* 2013;29(2).
21. Bohra N, Srivastava S, Bhatia M. Depression in women in Indian context. *Indian J Psychiatr.* 2015;57(7):239e245.

22. Ayaz SA, Mujahid S, Aatif S, Mukhtar M, Iftuquar S. Antiulcerogenic activity of *Caesalpinia pulcherrima* leaves. *Int J Pharm Res Allied Sci.* 2015;4(2):74-78.
23. Shah SK, Jhade DN, Chouksey R, Pharmacological Evaluation and Antifertility Activity of *Aloe Barbadensis* Linn on Female Wistar Rats, *Int. J. Phytomedicine*, 2017;9(2):253-260..
24. Christina L, Chichioco- Hernandez, Finella Marie G, Leonido. Weight- lowering effects of *Caesalpinia pulcherrima*, *Cassia fistula* and *Senna alata* leaf extracts. *J Medicinal Plants Res.* 2011;5(3):452-455.
25. Kandi P, Hayslett RL. Nicotine and 17 $\beta$ -estradiol produce an antidepressant-like effect in female ovariectomized rats. *Brain Res Bull.* 2011;84(3):224-228.
26. Swaab FD, Bao MA, Lucassen JP. The stress system in the human brain in depression and neurodegeneration. *Ageing: Res.Rev.* 2005;4(2):141-194.
27. Zhao D, Zheng L, Qi L, Wang S, Guan L, Xia Y, et al. Structural features and potent antidepressant effects of total sterols and beta-sitosterol extracted from *Sargassum horneri*. *Mar Drugs.* 2016; 14(7): 123.
28. Chuang CY, Shi YC, You HP, et al. Antidepressant effect of GABA-rich *Monascus* fermented product on forced swimming rat model. *J Agric Food Chem.* 2011;59(7):3027e3034.
29. Fisar Z, Hroudova J, Raboch J. Inhibition of monoamine oxidase activity by antidepressants and mood stabilizers. *Neuroendocrinol Lett.* 2010;31(5):42.
30. Lee CY, Cheng HM, Sim SM. Mulberry leaves protect rat from immobilization stress-induced inflammation. *Biofactors* 2007; 31(1):25–33.
31. Chutia D, Tyagi C.K., Bhuyan N.R, Shah SK, In vivo and in vitro Hyperglycaemic Screening of *Adenocalymma alliaceum* Miers, A Promising Herb, *Indian Journal of Pharmaceutical Education and Research*, 2022;56(4): S713-S724
32. Kavith AN, Naira N. Formulation and evaluation of the methanol extract of *Caesalpinia pulcherrima* leaves for its wound healing activity. *Asian J Pharm Res Health Care.* 2012;4(3):90-94.
33. Tyagi CK, Jhade D, Shah SK, Isolation, evaluation and characterization of isolated compounds from aqueous extract of *Cestrum nocturnum*, *Int J Phytomed*, 2017;8; 491-499.
34. Pal, S. Dutta, A. Sarkar Evaluation of CNS activities of ethanol extract of roots and rhizomes of *Cyperus rotundus* in mice, *Acta Pol Pharm - Drug Res.*, 66 (5) (2009), pp. 535-541
35. Onasanwo SA, Ilenre KO, Faborode SO. The impact of Kolaviron (a biflavonoid of *Garcinia kola* seed) on depression status in laboratory rodents: Roles of monoaminergic systems. *Ann Depress Anxiety*, 2015; 2(1):1042.
36. Cryan FJ, Valentino RJ, Lucki I. Assessing substrates underlying the behavioral effects of antidepressants using the modified rat forced swimming test. *Neurosci Biobehav Rev*, 2005;29:547–69.